

# Nizoral<sup>TRADEMARK</sup>

(ketoconazole) tablets

# over 5 million prescriptions world-wide.

## Oral medication in antifungal therapy:



Doctors and patients around the world are discovering the modern simplicity of Nizoral oral therapy. Common but often problematic fungal infections can now be treated effectively and elegantly.

Typically, in *Candida vaginitis*, 2 oral tablets once daily for 5 days is all it takes today to effectively cure the problem.



**JANSSEN**  
PHARMACEUTICA

*the drug discovery company*

#### Prescribing Information

**Presentation:** white, flat, half scored uncoated tablets marked "Janssen" on one side and K/200 on the reverse. Each tablet contains 200 mg ketoconazole. **Uses:** Nizoral is an orally active antimycotic for the treatment in adults of vaginal candidosis, superficial and systemic mycoses including dermatophyte and yeast infections of the skin, hair and nails, yeast infections of the mouth and G.I.-tract. Also maintenance treatment of systemic mycoses and chronic mucocutaneous candidosis and prophylaxis in "at risk" patients. In children: systemic mycoses and severe local infections where previous topical treatment has failed. **Side-effects, precautions, contra-indications:** contra-indicated in pregnancy. For maximal absorption Nizoral should be taken with meals. The use of agents which reduce gastric acidity (anti-cholinergic drugs, antacids, H<sub>2</sub>-blockers) should be avoided and, if indicated, such drugs should be taken not less than two hours after Nizoral. Nausea, skin rash, headache and pruritus may occasionally be observed. Alterations in liver function tests have occurred in patients on ketoconazole, these changes may be transient. Cases of hepatitis have been reported with an incidence of about 1 per 10,000 patients. Some of these may represent an idiosyncratic adverse reaction to the drug. This should be borne in mind in patients on long-term therapy. If a patient develops jaundice or any symptoms suggestive of hepatitis, treatment with ketoconazole should be stopped. Mild asymptomatic increases of liver enzyme levels, on the other hand, do not necessitate discontinuation of the treatment. Ketoconazole, when given together with cyclosporin A results in increased blood levels of cyclosporin A. It is important that blood levels of cyclosporin A are monitored if the two drugs are given concomitantly. Ketoconazole 200 mg once daily produces a transient decrease in plasma levels of testosterone. During long-term therapy at this dose, testosterone levels are usually not significantly different from controls. In rare instances, gynaecomastia has been reported.

# ARDUAN<sup>®</sup> vials and ampoules

## PIPECURONIUM BROMIDE

Arduan<sup>®</sup> is a new non-depolarizing muscle relaxant with a steroid structure.

Its main characteristics are:

- moderate duration of action
- circulatory stability
- effect can well be antagonised
- it has no histamine liberating and cumulative effect

Produced by Chemical Works of Gedeon Richter Ltd.

Exported by **medimpex**



Hungarian Trading Company for Pharmaceutical Products  
H-1808 Budapest telex 22 5477  
Phone 183-955-Hungary

In the EEC and some other countries the product will be marketed by ORGANON TEKNIKA



**arduan<sup>®</sup>**

# THE CARDIOPROTECTIVE CORONARY THERAPEUTIC AGENT ADALAT:

## THE MODERN COMPREHENSIVE DRUG THERAPY OF CORONARY HEART DISEASE.

**Composition:** 1 capsule of Adalat contains 10 mg of nifedipine. **Indications:** 1. Coronary heart disease: Chronic stable angina pectoris, angina at rest, including vasospastic angina pectoris (Prinzmetal's angina), variant angina) and unstable angina (crescendo pre-infarction angina), post-infarction angina (except for the first 8 days following the acute event). 2. Hypertension. **Contraindications:** Adalat must not be used at any time during pregnancy. No findings are as yet available as regards its use during the nursing period. Caution should be exercised if blood pressure is extremely low (severe hypotension: SBP < 90 mmHg). Cardiovascular shock. **Side effects:** Adverse reactions occur predominantly at the beginning of treatment and are often mild and transient: facial flush, heat sensation, headache. In isolated cases after high doses: nausea, dizziness, fatigue, skin reactions, paraesthesia, hypotensive reaction, palpitations and rise in pulse rate. Occasionally leg oedemas may occur as a result of vasodilation. Very rare: Gingival hyperplasia may develop during prolonged treatment, however, this is completely reversible after medication has been stopped; chest pain (possibly angina pectoris-like symptoms), in this case nifedipine should be discontinued, if a causal relationship is suspected. Caution must be exercised in patients who have to undergo regular dialysis and suffer from malignant hypertension and in irreversible renal failure associated with hypovolaemia, since blood pressure may fall as a result of vasodilation. Antihypertensive treatment with this drug requires regular medical check-ups. The response to drug intake may in individual cases be such that the ability to drive or operate machinery is impaired. These reactions are especially pronounced at the start of treatment and in the case of a change in medication and, moreover, if alcohol is consumed at the same time. **Interactions with other drugs:** Nifedipine/antihypertensives: Concomitant use may intensify the blood pressure-lowering effect of nifedipine. Nifedipine/beta-receptor blockers: Concomitant use requires careful monitoring of the patient, since the hypotensive effect may be enhanced; moreover, the development of heart failure has occasionally been reported. Nifedipine/cimetidine: The blood pressure-lowering effect may be intensified. **Presentation:** Adalat: Packages containing 30, 50 and 100 capsules of 10 mg of nifedipine. Co-No.: 1a. Bayer, Germany.



Bayer