

Nizoral^{TRADEMARK}

(ketoconazole) tablets

over 5 million
prescriptions
world-wide.

Oral medication
in antifungal therapy:



Doctors and patients around the world are discovering the modern simplicity of Nizoral oral therapy. Common but often problematic fungal infections can now be treated effectively and elegantly.

Typically, in *Candida* vaginitis, 2 oral tablets once daily for 5 days is all it takes today to effectively cure the problem.

 **JANSSEN**
PHARMACEUTICA

the drug discovery company

Prescribing information

Presentation: white, flat, half-scored uncoated tablets marked "Janssen" on one side and "K 200" on the reverse. Each tablet contains 200 mg ketoconazole. **Uses:** Nizoral is an orally active antimycotic for the treatment in adults of vaginal candidosis, superficial and systemic mycoses including dermatophyte and yeast infections of the skin, hair and nails, yeast infections of the mouth and G.I. tract. Also maintenance treatment of systemic mycoses and chronic mucocutaneous candidosis and prophylaxis in at-risk patients. In children, systemic mycoses and severe local infections where previous topical treatment has failed. **Side-effects, precautions, contra-indications:** contra-indicated in pregnancy. For maximal absorption Nizoral should be taken with meals. The use of agents which reduce gastric acidity (anti-cholinergic drugs, antacids, H₂-blockers) should be avoided and, if indicated, such drugs should be taken not less than two hours after Nizoral. Nausea, skin rash, headache and pruritus may occasionally be observed. Alterations in liver function tests have occurred in patients on ketoconazole; these changes may be transient. Cases of hepatitis have been reported with an incidence of about 1 per 10 000 patients. Some of these may represent an idiosyncratic adverse reaction to the drug. This should be borne in mind in patients on long-term therapy. If a patient develops jaundice or any symptoms suggestive of hepatitis, treatment with ketoconazole should be stopped. Mild asymptomatic increases of liver enzyme levels, on the other hand, do not necessitate discontinuation of the treatment. Ketoconazole, when given together with cyclosporin A, results in increased blood levels of cyclosporin A. It is important that blood levels of cyclosporin A are monitored if the two drugs are given concomitantly. Ketoconazole 200 mg once daily produces a transient decrease in plasma levels of testosterone. During long-term therapy at this dose, testosterone levels are usually not significantly different from controls. In rare instances, gynaecomastia has been reported.


ABC OF PLASTIC AND RECONSTRUCTIVE SURGERY

DAI DAVIES

Essential facts on plastic surgery

Spectacular advances in the technique of plastic and reconstructive surgery mean that surgeons can now reposition amputated limbs and restore function to lacerated hands; they also play a vital part in multidisciplinary teams treating patients with head and neck cancer. *ABC of Plastic and Reconstructive Surgery* brings the non-specialist up to date on these advances and explains what can be done about congenital abnormalities and the disfiguring effects of trauma or disease. Fully illustrated to show the surgical procedures used and the results that can be achieved, it includes chapters on the whole range of reconstructive problems

from burns to malignancies. A series from the *British Medical Journal*, now collected in one convenient volume with specially improved reproduction of the illustrations.



Price: Inland £6.50; Abroad £9.00; USA \$15.00
BMA members: Inland £6.00; Abroad £8.50; USA \$14.00
(Please quote membership number)
Prices include postage, by air abroad.
Payment must be enclosed with order.

BOOKS FROM THE BMJ

Order from The Publishing Manager, British Medical Journal, BMA House, Tavistock Square,
London WC1H 9JR or any leading bookseller

MEDICINE IN OLD AGE

Second Edition

A new look at old age . . .

The need to provide proper health care for elderly patients is an increasing challenge to modern medicine. This second, revised edition of *Medicine in Old Age* brings the general medical reader up to date on recent advances in geriatrics. It includes accounts of the management of dementia, gastrointestinal problems, anaemia, and urinary tract diseases as well as guidance on drug treatment and the specific problems of interpreting laboratory findings in the old. Chapters on rehabilitation and the role of the general practitioner and day hospitals—which have helped to convert mere existence into a full life for many elderly patients—complete a useful introduction to this important subject.



Price: Inland £6.95
Abroad £9.45/USA \$14.00
(Inland £6.45
Abroad £8.95/USA \$13.00
to BMA members. Please quote membership
number.)

Prices include postage, by air abroad

Payment must be enclosed with order

Order your copy now

From: The Publishing Manager
British Medical Journal
BMA House
Tavistock Square
London WC1H 9JR
or any leading bookseller

Now. Worldwide medical services for patients and families where specialized care and treatment is required.



These are unique, highly specialized services offered by International Health Systems through thirteen medical institutions located throughout the United States.

All are internationally recognized for their excellence in treating cardiac problems, renal diseases, endocrine problems, high-risk obstetrical care, neurological disorders, birth defects,

cancer, strokes and accidents (for which physical medicine and rehabilitation are provided), children's disorders, and many other medical problems.

International Health Systems accommodates the patient's and family's special needs, while in the U.S., and provides a continuation of care after the patient returns home.

The IHS affiliated medical centers include: University of California at San Diego & Los Angeles, University of Florida, Emory University, University of Hawaii, Northwestern University Medical School, Washington University, Duke University Hospital, Baylor College of Medicine, University of Texas at San Antonio, University of Utah, University of Washington and The George Washington University.

INTERNATIONAL HEALTH SYSTEMS

For more information mail to:

International Health Systems, Inc.
No. 3 Berkeley Square
London, W1X 5HG Telephone: 01-491-3611
Telex: 267383 Facsimile: 01-491-4811
Or
3773 Cherry Creek North Drive
Denver, CO 80209 U.S.A.
Telephone: 303-399-2105 Telex: 820107



Dear Sirs:

☐ Please Send Your Brochure And Application On Services Available

☐ Please Have A Representative Contact Me

(Please Print)

Name/Title

Institution/Organization

Address


City/State/Country



THE CARDIOPROTECTIVE CORONARY THERAPEUTIC AGENT ADALAT®

THE MODERN COMPREHENSIVE DRUG THERAPY OF CORONARY HEART DISEASE.

Composition: 1 capsule of Adalat contains 10 mg of nifedipine. **Indications:** 1. Coronary heart disease: Chronic stable angina pectoris, angina at rest, including vasospastic angina pectoris (Prinzmetal's angina, variant angina) and unstable angina (crescendo pre-infarction angina), post-infarction angina (except for the first 8 days following the acute event). 2. Hypertension. **Contraindications:** Adalat must not be used at any time during pregnancy. No findings are as yet available as regards its use during the nursing period. **Caution should be exercised if blood pressure is extremely low (severe hypotension: SBP < 90 mmHg). Cardiovascular shock. Side effects:** Adverse reactions occur predominantly at the beginning of treatment and are often mild and transient: facial flush, heat sensation, headache. In isolated cases after high doses: nausea, dizziness, fatigue, skin reactions, paraesthesia, hypotensive reaction, palpitations and rise in pulse rate. Occasionally leg oedemas may occur as a result of vasodilation. Very rare: Gingival hyperplasia may develop during prolonged treatment, however, this is completely reversible after medication has been stopped; chest pain (possibly angina pectoris-like symptoms), in this case nifedipine should be discontinued, if a causal relationship is suspected. **Caution must be exercised in patients who have to undergo regular dialysis and suffer from malignant hypertension and in irreversible renal failure associated with hypovolaemia, since blood pressure may fall as a result of vasodilation. Antihypertensive treatment with this drug requires regular medical check-ups. The response to drug intake may in individual cases be such that the ability to drive or operate machinery is impaired. These reactions are especially pronounced at the start of treatment and in the case of a change in medication and, moreover, if alcohol is consumed at the same time. Interactions with other drugs:** Nifedipine/antihypertensives: Concomitant use may intensify the blood pressure-lowering effect of nifedipine. Nifedipine/beta-receptor blockers: Concomitant use requires careful monitoring of the patient, since the hypotensive effect may be enhanced; moreover, the development of heart failure has occasionally been reported. Nifedipine/cimetidine: The blood pressure-lowering effect may be intensified. **Presentation:** Adalat: Packages containing 30, 50 and 100 capsules of 10 mg of nifedipine. Co-No.: 1a. Bayer, Germany.



Bayer