

Nizoral

TRADEMARK

(ketoconazole)

tablets

over 5 million prescriptions world-wide.

Oral medication in antifungal therapy:



Doctors and patients around the world are discovering the modern simplicity of Nizoral oral therapy. Common but often problematic fungal infections can now be treated effectively and elegantly.

Typically, in *Candida* vaginitis, 2 oral tablets once daily for 5 days is all it takes today to effectively cure the problem.



JANSSEN
PHARMACEUTICA


the drug discovery company

Prescribing Information

Presentation: white, flat, half scored uncoated tablets marked "Janssen" on one side and K/200 on the reverse. Each tablet contains 200 mg ketoconazole. **Uses:** Nizoral is an orally active antimycotic for the treatment in adults of vaginal candidosis, superficial and systemic mycoses including dermatophyte and yeast infections of the skin, hair and nails, yeast infections of the mouth and G.I.-tract. Also maintenance treatment of systemic mycoses and chronic mucocutaneous candidosis and prophylaxis in "at risk" patients. In children: systemic mycoses and severe local infections where previous topical treatment has failed. **Side-effects, precautions, contra-indications:** contra-indicated in pregnancy. For maximal absorption Nizoral should be taken with meals. The use of agents which reduce gastric acidity (anti-cholinergic drugs, antacids, H₂-blockers) should be avoided and, if indicated, such drugs should be taken not less than two hours after Nizoral. Nausea, skin rash, headache and pruritus may occasionally be observed. Alterations in liver function tests have occurred in patients on ketoconazole, these changes may be transient. Cases of hepatitis have been reported with an incidence of about 1 per 10,000 patients. Some of these may represent an idiosyncratic adverse reaction to the drug. This should be borne in mind in patients on long-term therapy. If a patient develops jaundice or any symptoms suggestive of hepatitis, treatment with ketoconazole should be stopped. Mild asymptomatic increases of liver enzyme levels, on the other hand, do not necessitate discontinuation of the treatment. Ketoconazole, when given together with cyclosporin A, results in increased blood levels of cyclosporin A. It is important that blood levels of cyclosporin A are monitored if the two drugs are given concomitantly. Ketoconazole 200 mg once daily produces a transient decrease in plasma levels of testosterone. During long-term therapy at this dose, testosterone levels are usually not significantly different from controls. In rare instances, gynaecomastia has been reported.

THE KEYNES PRESS

Limited editions of
medical classics,
published under an imprint
named after Sir Geoffrey Keynes



	UK	Abroad	USA		UK	Abroad	USA
<i>The Portraiture of William Harvey</i> by Sir Geoffrey Keynes	£40.00	£46.00	\$64.50	<i>Platt versus Pickering</i>	£45.00	£51.00	\$71.50
<i>Instincts of the Herd in Peace and War</i> , by William Trotter	£45.00	£52.00	\$73.00	<i>Notes on the Composition of Scientific Papers</i> , by Sir Clifford Allbutt	£35.00	£39.50	\$55.50
<i>A Varied Life:</i> <i>Memoirs of a Swansea GP</i> , the diaries of Urban Marks	£35.00	£37.50	\$52.50	<i>The Anatomy of Courage</i> , by Lord Moran	£35.00	£41.00	\$57.50

Order from the Publishing Manager (Keynes Press), BMA House, Tavistock Square, London WC1H 9JR.
All prices include registered postage, by air abroad. Please make your cheque payable to the *British Medical Journal*.

Rocephin


ceftriaxone

Third-Generation

24-hour bactericidal power

Third-Generation cephalosporin

«Rocephin»: 24-hour bactericidal power



Comparison of concentrations in plasma and in extracellular fluid (skin blisters) after a single dose of 0.5 g ceftriaxone or 1 g cefotaxime (from Wise and Andrews¹).

«Rocephin» longer

1. Wise R, Andrews B. A comparison of the bactericidal activity and tissue penetration of ceftriaxone, cefotaxime and cefotaxime. *Eur J Clin Microbiol* 2, 505-508 (1983).



F. Hoffmann-La Roche & Co.
Limited Company, Basle, Switzerland

Full details on composition, indications, contraindications, dosage and precautions are available on request.