

Nizoral^{TRADEMARK}

(ketoconazole) tablets

over 5 million
prescriptions
world-wide.

**Oral medication
in antifungal therapy:**



Doctors and patients around the world are discovering the modern simplicity of Nizoral oral therapy. Common but often problematic fungal infections can now be treated effectively and elegantly.

Typically, in *Candida* vaginitis, 2 oral tablets once daily for 5 days is all it takes today to effectively cure the problem.



**JANSSEN
PHARMACEUTICA**

the drug discovery company

Prescribing Information:

Presentation: white, flat, half scored uncoated tablets marked "Janssen" on one side and K/200 on the reverse. Each tablet contains 200 mg ketoconazole. **Uses:** Nizoral is an orally active antimycotic for the treatment in adults of vaginal candidosis, superficial and systemic mycoses including dermatophyte and yeast infections of the skin, hair and nails, yeast infections of the mouth and G.I. tract. Also maintenance treatment of systemic mycoses and chronic mucocutaneous candidosis and prophylaxis in "at risk" patients. In children: systemic mycoses and severe local infections where previous topical treatment has failed. **Side-effects, precautions, contra-indications:** contra-indicated in pregnancy. For maximal absorption Nizoral should be taken with meals. The use of agents which reduce gastric acidity (anti-cholinergic drugs, antacids, H₂-blockers) should be avoided and, if indicated, such drugs should be taken not less than two hours after Nizoral. Nausea, skin rash, headache and pruritus may occasionally be observed. Alterations in liver function tests have occurred in patients on ketoconazole, these changes may be transient. Cases of hepatitis have been reported with an incidence of about 1 per 10,000 patients. Some of these may represent an idiosyncratic adverse reaction to the drug. This should be borne in mind in patients on long-term therapy. If a patient develops jaundice or any symptoms suggestive of hepatitis, treatment with ketoconazole should be stopped. Mild asymptomatic increases of liver enzyme levels, on the other hand, do not necessitate discontinuation of the treatment. Ketoconazole, when given together with cyclosporin A results in increased blood levels of cyclosporin A. It is important that blood levels of cyclosporin A are monitored if the two drugs are given concomitantly. Ketoconazole 200 mg once daily produces a transient decrease in plasma levels of testosterone. During long-term therapy at this dose, testosterone levels are usually not significantly different from controls. In rare instances, gynecomastia has been reported.

JUMEX[®]

Tablets

(Selegilinum hydrochloricum) Selective MAO-B Inhibitor

...A STEP FORWARD IN THE TREATMENT OF PARKINSONISM



JUMEX[®]

- develops its effect only in conjunction with levodopa preparations
- doses of levedopa can be reduced while maintaining the therapeutic effect
- prolongs the effective period of levodopa
- is applicable in all phases of Parkinsonism
- on-off periods decline in frequency and extent
- improves quality of life by diminishing side-effects
- improves motor activity
- milder end-of-dose akinesia
- simple dosage:
1-2 tablets (5-10 mg) daily
- composition:
5 mg Selegilinum
hydrochloricum (L-deprenyl)
per tablet

Manufacturer
CHINOIN Pharmaceutical
and Chemical Works — Budapest



Exported by:
MEDIMPEX • HUNGARY
H - 1808 BUDAPEST, PHONE: 183-955, TX: 22-5477

GOOD DRUGS DON'T GROW ON TREES

*In the prehistoric days of medicine
some drugs could be found
growing on trees.
Today that's a bit different.*

*The therapeutic drugs of our times
are being unmasked from nature
rather more scientifically.
By thorough bio-medical research
into the fundamental processes of life.
By studying exactly why and how
these processes sometimes go wrong.*

*And so nature shows man where and how
he can intervene and defend himself.
With precision drugs that will selectively
and safely correct what went wrong.
Today's drugs grow from man's scientific
knowledge of nature —
but no longer on trees.*