

Nizoral^{TRADEMARK}

(ketoconazole) tablets

over 5 million prescriptions world-wide.

Oral medication in antifungal therapy:



Doctors and patients around the world are discovering the modern simplicity of Nizoral oral therapy. Common but often problematic fungal infections can now be treated effectively *and elegantly*.

Typically, in *Candida* vaginitis, 2 oral tablets once daily for 5 days is all it takes today to effectively cure the problem.



JANSSEN
PHARMACEUTICA

the drug discovery company

Prescribing Information

Presentation: white, flat, half scored uncoated tablets marked "Janssen" on one side and K/200 on the reverse. Each tablet contains 200 mg ketoconazole. **Uses:** Nizoral is an orally active antimycotic for the treatment in adults of vaginal candidosis, superficial and systemic mycoses including dermatophyte and yeast infections of the skin, hair and nails, yeast infections of the mouth and G.I.-tract. Also maintenance treatment of systemic mycoses and chronic mucocutaneous candidosis and prophylaxis in "at risk" patients. In children: systemic mycoses and severe local infections where previous topical treatment has failed. **Side-effects, precautions, contra-indications:** contra-indicated in pregnancy. For maximal absorption Nizoral should be taken with meals. The use of agents which reduce gastric acidity (anti-cholinergic drugs, antacids, H₂-blockers) should be avoided and, if indicated, such drugs should be taken not less than two hours after Nizoral. Nausea, skin rash, headache and pruritus may occasionally be observed. Alterations in liver function tests have occurred in patients on ketoconazole, these changes may be transient. Cases of hepatitis have been reported with an incidence of about 1 per 10,000 patients. Some of these may represent an idiosyncratic adverse reaction to the drug. This should be borne in mind in patients on long-term therapy. If a patient develops jaundice or any symptoms suggestive of hepatitis, treatment with ketoconazole should be stopped. Mild asymptomatic increases of liver enzyme levels, on the other hand, do not necessitate discontinuation of the treatment. Ketoconazole, when given together with cyclosporin A results in increased blood levels of cyclosporin A. It is important that blood levels of cyclosporin A are monitored if the two drugs are given concomitantly. Ketoconazole 200 mg once daily produces a transient decrease in plasma levels of testosterone. During long-term therapy at this dose, testosterone levels are usually not significantly different from controls. In rare instances, gynaecomastia has been reported.

EPIDEMIOLOGY FOR THE UNINITIATED

GEOFFREY ROSE, D J P BARKER


SECOND EDITION

Get a working knowledge of epidemiology

No one would expect to understand a disease without knowledge of its clinical findings and pathology, but a surprising number of doctors remain ignorant of another important aspect – the study of disease in relation to populations. Epidemiology has its own techniques of data collection and interpretation and its necessary jargon of technical terms, and in *Epidemiology for the Uninitiated* Professors Geoffrey Rose and David Barker guide the novice expertly through the theory and practical pitfalls. The second edition of this popular *BMJ* handbook has been revised to include further details of epidemiological methods and some of their more dramatic applications, such as the investigations on the Spanish cooking oil epidemic, and AIDS.

Price: Inland £3.95; Abroad £5.50;
USA \$8.50
BMA members; Inland £3.45; Abroad £5.00;
USA \$7.50

(Please quote membership number)
Prices include postage, by air abroad.
Payment must be enclosed with order.



Reviews of the first edition

"... all physicians [should] ensure that their hospital library has this book, and that they themselves read it."

Med J Aust 1981; 1: 665

"A reading of this book ... should provide a sound basis in epidemiological techniques for any intending researcher."

NZ Med J 1980; 91: 73

BOOKS FROM THE BMJ

Order from The Publishing Manager, British Medical Journal, BMA House, Tavistock Square,
London WC1H 9JR or any leading bookseller

Trade Mark

daily cephalosporin

«Roceph» longer



ROCHE

Full details on composition, actions, contraindications, dosage and precautions are available on request.