

Nizoral^{TRADEMARK}

(ketoconazole) tablets

over 5 million
prescriptions
world-wide.

Oral medication
in antifungal therapy:



Doctors and patients around the world are discovering the modern simplicity of Nizoral oral therapy. Common but often problematic fungal infections can now be treated effectively and elegantly.

Typically, in *Candida* vaginitis, 2 oral tablets once daily for 5 days is all it takes today to effectively cure the problem.



JANSSEN
PHARMACEUTICA

the drug discovery company


Prescribing Information

Presentation: white, flat, half scored uncoated tablets marked "Janssen" on one side and K/200 on the reverse. Each tablet contains 200 mg ketoconazole. **Uses:** Nizoral is an orally active antimycotic for the treatment in adults of vaginal candidosis, superficial and systemic mycoses including dermatophyte and yeast infections of the skin, hair and nails, yeast infections of the mouth and G.I. tract. Also maintenance treatment of systemic mycoses and chronic mucocutaneous candidosis and prophylaxis in "at risk" patients. In children: systemic mycoses and severe local infections where previous topical treatment has failed. **Side-effects, precautions, contraindications:** contraindicated in pregnancy. For maximal absorption Nizoral should be taken with meals. The use of agents which reduce gastric acidity (anti-cholinergic drugs, antacids, H₂-blockers) should be avoided and, if indicated, such drugs should be taken not less than two hours after Nizoral. Nausea, skin rash, headache and pruritus may occasionally be observed. Alterations in liver function tests have occurred in patients on ketoconazole, these changes may be transient. Cases of hepatitis have been reported with an incidence of about 1 per 10,000 patients. Some of these may represent an idiosyncratic adverse reaction to the drug. This should be borne in mind in patients on long-term therapy. If a patient develops jaundice or any symptoms suggestive of hepatitis, treatment with ketoconazole should be stopped. Mild asymptomatic increases of liver enzyme levels, on the other hand, do not necessitate discontinuation of the treatment. Ketoconazole, when given together with cyclosporin A results in increased blood levels of cyclosporin A. It is important that blood levels of cyclosporin A are monitored if the two drugs are given concomitantly. Ketoconazole 200 mg once daily produces a transient decrease in plasma levels of testosterone. During long-term therapy at this dose, testosterone levels are usually not significantly different from controls. In rare instances, gynaecomastia has been reported.

Tobanur®

non-cardioselective
beta-blocker

tablets
(CLORANOLOL)



Cloranolol is a non-cardio-selective beta-adrenergic receptor blocking agent. It is 3 to 10 times more active than propranolol. It does not possess any sympathomimetic action. It exerts a membrane-stabilizing "channel-like" effect. It is highly effective in the treatment of hypertension, angina pectoris, different irregularities of the cardiac rhythm, essential circulatory hyperkinesia, hypertrophic obstructive cardiomyopathy and hyperthyroidism.

Produced by
**CHEMICAL WORKS of
GEDEON RICHTER Ltd.**
Budapest-Hungary

Exported by
MEDIMPEX
Budapest-Hungary

WHAT'S A GOOD DOCTOR WITHOUT GOOD DRUGS ?



Good doctors treat patients, not just diseases or symptoms. Diagnosis is their key to selecting the most appropriate treatment, always weighing risks against benefits.

And when it comes to drug treatment, they like to rely on the best products from innovative drug research.

They know that ever more effective and safer drugs are indispensable to progress in medicine.

JANSSEN
PHARMACEUTICA
the discovery company