

Nizoral^{TRADEMARK}

(ketoconazole) tablets

over 5 million prescriptions world-wide.

Oral medication in antifungal therapy:



Doctors and patients around the world are discovering the modern simplicity of Nizoral oral therapy. Common but often problematic fungal infections can now be treated effectively *and elegantly*.

Typically, in *Candida* vaginitis, 2 oral tablets once daily for 5 days is all it takes today to effectively cure the problem.



**JANSSEN
PHARMACEUTICA**

the drug discovery company

Prescribing Information

Presentation: white, flat, half scored uncoated tablets marked "Janssen" on one side and K/200 on the reverse. Each tablet contains 200 mg ketoconazole. **Uses:** Nizoral is an orally active antimycotic for the treatment in adults of vaginal candidosis, superficial and systemic mycoses including dermatophyte and yeast infections of the skin, hair and nails, yeast infections of the mouth and G.I.-tract. Also maintenance treatment of systemic mycoses and chronic mucocutaneous candidosis and prophylaxis in "at risk" patients. In children: systemic mycoses and severe local infections where previous topical treatment has failed. **Side-effects, precautions, contra-indications:** contra-indicated in pregnancy. For maximal absorption Nizoral should be taken with meals. The use of agents which reduce gastric acidity (anti-cholinergic drugs, antacids, H₂-blockers) should be avoided and, if indicated, such drugs should be taken not less than two hours after Nizoral. Nausea, skin rash, headache and pruritus may occasionally be observed. Alterations in liver function tests have occurred in patients on ketoconazole, these changes may be transient. Cases of hepatitis have been reported with an incidence of about 1 per 10,000 patients. Some of these may represent an idiosyncratic adverse reaction to the drug. This should be borne in mind in patients on long-term therapy. If a patient develops jaundice or any symptoms suggestive of hepatitis, treatment with ketoconazole should be stopped. Mild asymptomatic increases of liver enzyme levels, on the other hand, do not necessitate discontinuation of the treatment. Ketoconazole, when given together with cyclosporin A results in increased blood levels of cyclosporin A. It is important that blood levels of cyclosporin A are monitored if the two drugs are given concomitantly. Ketoconazole 200 mg once daily produces a transient decrease in plasma levels of testosterone. During long-term therapy at this dose, testosterone levels are usually not significantly different from controls. In rare instances, gynaecomastia has been reported.

ARDUAN[®] vials and ampoules

PIPECURONIUM BROMIDE

Arduan[®] is a new non-depolarizing muscle relaxant with a steroid structure.

Its main characteristics are:

- moderate duration of action
- circulatory stability
- effect can well be antagonised
- it has no histamine liberating and cumulative effect

Produced by Chemical Works of Gedeon Richter Ltd.

Exported by **medimpex**

Hungarian Trading Company for Pharmaceutical Products
H-1808 Budapest telex 22 5477
Phone 183-955-Hungary



In the EEC and some other countries the product will be marketed by ORGANON TEKNIKA



GOOD DRUGS DON'T GROW ON TREES

*In the prehistoric days of medicine
some drugs could be found
growing on trees.
Today that's a bit different.*

*The therapeutic drugs of our times
are being unmasked from nature
rather more scientifically.*

*By thorough bio-medical research
into the fundamental processes of life.
By studying exactly why and how
these processes sometimes go wrong.*

*And so nature shows man where and how
he can intervene and defend himself.
With precision drugs that will selectively
and safely correct what went wrong.
Today's drugs grow from man's scientific
knowledge of nature —
but no longer on trees.*