

In vaginal candidosis:

TRADEMARK

The elegant way to treat an inelegant problem ketoconazole





PA helps in the early detection of recurrent carcinomas. And early treatment can improve the prognosis and quality of life for your patient. Rises in the post-treatment levels of serum TPA often precede the clinical symptoms of recurrent carcinomas by as much as six months.* Serum TPA (Tissue Polypeptide Antigen) can be measured quickly and easily using the Prolifigen® TPA IRMA test.

*Documented for carcinomas of the bladder, breast, lung and genital tract.

The illustration depicts the little Dutch boy Hans Brinker saving his home-town of Haarlem from flooding by sticking his finger into a hole in the dyke. The story was written by Mary Mapes Dodge in 1873.

This legend about the fair-haired hero of Haarlem is one of the world's best known. As with all good fairy-tales, it has a moral: "Don't wait until too late."


Serum TPA measurements: For early and efficient detection of recurrent malignant disease.



Box 20045, 5-16120 Bromma, Sweden. Tel. Int +468988330.

Just published

One revolution leads to another



The extraordinary technical developments in molecular biology over the past few years, and the equally rapid advances in understanding of cell biology, will almost certainly result in far reaching changes in medical research and practice. In this collection of articles experts in molecular and cell biology provide the background information to give clinicians an insight into the way in which the medical sciences may be moving over the next few years and into the exciting possibilities opening up for the treatment of genetic disorders, cancer, and the common illnesses of Western society such as degenerative vascular disease and diabetes.

Price: Inland £5.95; Abroad £7.50; USA \$12.00


BMA members: Inland £5.45; Abroad £7.00; USA \$11.00

including postage, by air abroad

Payment must be enclosed with order.

JUST PUBLISHED

Non-specialists overwhelmed by too many systematic descriptions of individual disease will welcome the new approach of ABC of Dermatology. Here the history, clinical appearance, and pathology of a few common key conditions are discussed and then used as a basis for comparison with other skin diseases. Also dealing in detail with allergic reactions, autoimmunity, acne, and infections, this handbook is fully illustrated in colour, providing a clear guide to assessment and treatment as well as an understanding of the histological changes underlying the clinical presentations.






Inland £10.95; Abroad £13.00; USA \$22.00 BMA members: Inland £9.95; Abroad £12.00; USA \$20.00 including postage by air abroad

Please enclose payment with order or send us full details of your ACCESS or VISA credit card.

ORDER FROM
BRITISH MEDICAL JOURNAL
PO Box 295, London WC1H 9TE
or any leading medical bookseller.

without clouding his head



Hismanal

The antihistamine of exceptional strength from the first to the last day of therapy.



Prescribing information

Prescribing information
Uses: Hismanal' is a potent and non-sedative antihistamine indicated for the treatment of seasonal and perennial allergic rhinitis. allergic conjunctivitis, chronic urticaria and other allergic conditions.

Dosage and administration: adults and children over 12 years: 10 mg daily. Children 6-12 years: 5 mg once daily. Children younger than 6 years. 2 mg per 10 kg daily. Hismanal should be taken on an empty stomach. Precautions: Hismanal should be used in pregnant women only when, in the judgement of the physician, the potential benefits outweigh the possible hazards. Adverse reactions: weight gain may occur during prolonged treatment. Overdose: in case of overdose, gastric lavage should be followed by close observation and ECG monitoring as arrhythmias have occasionally occurred after intake of doses in excess of 200 mg.

Full prescribing information available on request