


In vaginal candidosis:

TRADEMARK

ketoconazole

*The elegant way
to treat an inelegant problem*



2 oral tablets
once daily for **5** days
is all it takes today
to cure vaginal candidosis

...gastric
...not less than
...of cyclosporin A. It is
...side effects: nausea, skin rash,
...have occurred in patients on ketoconazole;
...reported with an incidence of about 1 per 10,000 patients. Some of
...to the drug. This should be borne in mind in patients on long-term therapy.
...suggestive of hepatitis, treatment with ketoconazole should be stopped. Mild
...on the other hand, do not necessitate discontinuation of the treatment.

References:

Tooley, *et al.*: The Practitioner **229**, 655 (1985)
Benussi, *et al.*: Curr. Ther. Res. **31**(4), 511 (1982)



JANSSEN
PHARMACEUTICA

B-2340 Beerse, Belgium

world leader in antimycotic research

IN LIFE-THREATENING INFECTIONS...

◀ROCEPHIN▶



...THE TREATMENT FOR EMERGENCIES

FOR EXAMPLE: SEPSIS... PATHOGENS ERADICATED IN 96.3% OF ALL CASES

Reference
Data on file, F. Hoffmann-La Roche & Co. Limited Company, Basle, Switzerland.

Full details on composition, indications, contraindications, side effects, dosage and precautions are available on request.



F. HOFFMANN-LA ROCHE & CO. LIMITED COMPANY, BASLE, SWITZERLAND



no skin allergy
without clouding
his head

Hismanal TRADEMARK
astemizole

**The antihistamine of exceptional strength from the first
to the last day of therapy.**

 **JANSSEN**
PHARMACEUTICAL
2340 Beerse, Belgium

Prescribing information

Uses: Hismanal[®] is a potent and non-sedative antihistamine indicated for the treatment of seasonal and perennial allergic rhinitis, allergic conjunctivitis, chronic urticaria and other allergic conditions.
Dosage and administration: adults and children over 12 years: 10 mg daily. Children 6-12 years: 5 mg once daily. Children younger than 6 years: 2 mg per 10 kg daily. Hismanal should be taken on an empty stomach. **Precautions:** Hismanal should be used in pregnant women only when, in the judgement of the physician, the potential benefits outweigh the possible hazards. **Adverse reactions:** weight gain may occur during prolonged treatment. **Overdose:** in case of overdose, gastric lavage should be followed by close observation and ECG monitoring as arrhythmias have occasionally occurred after intake of doses in excess of 200 mg.

Full prescribing information available on request

Trademark