


In vaginal candidosis:

Nizoral[®] TRADEMARK

ketoconazole

ketoconazole

*The elegant way
to treat an inelegant problem*



2 oral tablets
once daily for **5** days
is all it takes today
to cure vaginal candidosis

each side and 1000 on the reverse. Each tablet contains ketoconazole 200 mg. Dosage: 200 mg once daily for 5 days. For maximal absorption Nizoral should be taken on an empty stomach. **Precautions:** the use of agents which reduce gastric acidity should be avoided and, if indicated, such drugs should be taken not less than 2 hours after Nizoral. Concomitant use of cyclosporin A results in increased blood levels of cyclosporin A. It is recommended that if the two drugs are given concomitantly, **Side-effects:** nausea, skin rash, headache, dizziness, taste perversion have been observed. Alterations in liver function tests have occurred in patients on ketoconazole; however, cases of hepatitis have been reported with an incidence of about 1 per 10,000 patients. Some of these may represent an idiosyncratic adverse reaction to the drug. This should be borne in mind in patients on long-term therapy. If jaundice or any symptoms suggestive of hepatitis, treatment with ketoconazole should be stopped. Mild elevations of liver enzyme levels, on the other hand, do not necessitate discontinuation of the treatment.

Information available on request.

References:

References:

Tooley, *et al.*: The Practitioner **229**, 655 (1985)
Benussi, *et al.*: Curr. Ther. Res. **31**(4), 511 (1982)



JANSSEN
PHARMACEUTICA

B-2340 Beerse, Belgium


world leader in antimycotic research

CLINICAL EFFICACY PROVEN WORLDWIDE

«ROCEPHIN»

24-HOUR BACTERICIDAL POWER

FOR EXAMPLE: CLINICAL SUCCESS RATE
IN MAJOR SERIOUS INFECTIONS




Reference
Data on file, F. Hoffmann-La Roche & Co. Limited Company, Basle, Switzerland.

Full details on composition, indications, contraindications, side effects, dosage and precautions are available on request.



F. HOFFMANN-LA ROCHE & CO. LIMITED COMPANY, BASLE, SWITZERLAND

Clearing his skin allergy
without clouding
his head



Hismanal TRADEMARK
astemizole

The antihistamine of exceptional strength from the first
to the last day of therapy.

 **JANSSEN**
PHARMACEUTICAL
2340 Beerse, Belgium

Prescribing information

Uses: Hismanal[®] is a potent and non-sedative antihistamine indicated for the treatment of seasonal and perennial allergic rhinitis, allergic conjunctivitis, chronic urticaria and other allergic conditions.
Dosage and administration: adults and children over 12 years: 10 mg daily. Children 6-12 years: 5 mg once daily. Children younger than 6 years: 2 mg per 10 kg daily. Hismanal should be taken on an empty stomach. **Precautions:** Hismanal should be used in pregnant women only when, in the judgement of the physician, the potential benefits outweigh the possible hazards. **Adverse reactions:** weight gain may occur during prolonged treatment. **Overdose:** in case of overdose, gastric lavage should be followed by close observation and ECG monitoring as arrhythmias have occasionally occurred after intake of doses in excess of 200 mg.

Full prescribing information available on request

Trademark