
In vaginal candidosis:

TRADEMARK

ketoconazole The elegant way to treat an inelegant problem




Second Edition

ABC OF HYPERTENSION

A short guide to a complex and rapidly developing topic, the second, revised edition of ABC of Hypertension reviews up to date information from all recent clinical trials on hypertension and includes the latest recommendations of the British Hypertension Society. With its comprehensive survey of drug treatment – including advice on treating hypertension in pregnancy and in the elderly – this is an invaluable source of facts and guidance for practising doctors and medical students.

Chapters include


- Blood pressure measurement in infancy and childhood
- Drug treatment
- Blood pressure reduction in the elderly
- Hypertension in pregnancy
- Detecting hypertensive patients
- Management of hypertension in general practice



Price: Inland £4.95; Abroad £6.50; USA\$10.50 BMA members: Inland £4.45; Abroad £6.00; USA\$9.50 including postage, by air abroad

Payment must be enclosed with order

ABC update – three new editions




Just published


As any general practitioner knows, diseases of the ear, nose and throat are some of the most common complaints that bring patients to the surgery – children suffering misery and discomfort and adults having to take time off work. In the second, revised edition of ABC of Ear, Nose and Throat Harold Ludman gives busy clinicians expert, up to date advice on assessing and managing these conditions.

Price: Inland £5.95; Abroad £7.50; USA\$14.00 BMA members: Inland £5.45; Abroad £7.00; USA\$13.00

One of today's most pressing medical and social problems is comprehensively reviewed in the ABC of Alcohol. With an extended introduction by Richard Smith, assistant editor of the BMJ, this second, revised and expanded edition brings readers up to date on current approaches to the diagnosis and treatment of alcoholism. Essential reading for all doctors who need to understand the appalling difficulties that this condition creates for patients and their families – and indeed for doctors themselves.

Price: Inland £5.95; Abroad £7.50; USA\$13.00 BMA members: Inland £5.45; Abroad £7.00; USA\$12.00





Many advances in the treatment of diabetes have been made in the past decade: self measurement of blood glucose, intravenous infusions and intramuscular insulin treatment for diabetic emergencies, continuous subcutaneous insulin infusion techniques, and new tests for autonomic function. The outlook for diabetic pregnancy has been transformed; photocoagulation for retinopathy is reducing blindness; and at last diabetic renal disease is receiving some attention. Dr Peter Watkin's articles set these advances in their clinical context, providing a practical guide to the management of diabetes for non-specialist doctors and nurses. There have already been several more developments both in understanding the causes of diabetes and improving its management, and this second, revised, edition incorporates the most important of these latest innovations.

Price: Inland £4.95; Abroad £6.50; USA\$11.00 BMA members: Inland £4.45; Abroad £6.00; USA\$10.00


All prices include postage by air abroad

ORDER FORM

The Publishing Manager, BRITISH MEDICAL JOURNAL, P.O. Box 295, London WC1H 9TE, England or any leading medical bookseller.

London WCIH 91E, England or any leading medical bookseller.	
Please send me the following books:	
	☐ I encloseBMA Membership No
	Please make cheques payable to to British Medical Journal or debit my credit card (please tick box)
	Access Visa American Express
Name	Card No.
Address	Card expiry date
	Signature
	Please send me a book catalogue

Mark your hypertensive patient



Hypertension is one of the most important risk factors for the development of cardiovascular disorders.

However, not only the lowering of the blood pressure should be considered during the treatment. For many antihypertensive agents have a negative effect on

the plasma lipids. It should be prevented that the removal of one risk factor (hypertension) creates another risk factor (high cholesterol level).


The beta blocker Visken® (pindolol) takes an

V I S K E N° THE LIPID PROOF BETA BLOCKER

exceptional place because it does not have an unfavourable effect on the plasma lipid level. It is advisable to take this characteristic into consideration when choosing a therapy. Because your choice now also determines the risks your patient runs in the future.

VISKEN® B-adrenoceptor blocking agent with intrinsic sympathomimetic activity (ISA). Presentations: pindolol. Tablets 1, 2.5, 5, 10 or 15 mg. Retard tablet 20 mg. Oral solution 5 mg/ml. Ampoules of 0.4 mg/2 ml. Indications: hypertension: angina pectoris (prevention); certain types of arrhythmias (see full product information). Contraindications: bronchial asthma, digitalis-resistant cardiac failure, cor pulmonale, marked bradycardia, 2nd and 3rd degree AV-block. Precautions: incipient or manifest heart failure (digitalization required), tendency to bronchospasm in non-asthmatic chronic obstructive lung disease; general anaesthesia; interruption of treatment should be gradual; in phaeochromocytoma combine with or-blocker; pre-existing peripheral vascular disease; severe renal failure; in patients receiving antidiabetic therapy; when breast-feeding; when driving vehicles or operating machinery. Interactions: avoid concomitant parenteral use of calcium channel blockers. Side effects: tiredness, dizziness, nausea, headache, sleep disturbances; rarely skin reactions and psychic symptoms. Packs: country specific.

his skin allergy without clouding his head



Hismanal

The antihistamine of exceptional strength from the first to the last day of therapy.



Uses: Hismanal' is a potent and non-sedative antihistamine indicated for the treatment of seasonal and perennial allergic rhinitis. allergic conjunctivitis, chronic urticana and other allergic conditions.

Dosage and administration: adults and children over 12 years: 10 mg daily. Children 6-12 years: 5 mg once daily. Children younger than 6 years: 2 mg per 10 kg daily. Hismanal should be taken on an empty stomach. Precautions: Hismanal should be used in pregnant women only when, in the judgement of the physician, the potential benefits outweigh the possible hazards. Adverse reactions: weight gain may occur during prolonged treatment. Overdose: in case of overdose, gastric lavage should be followed by close observation and ECG monitoring as arrhythmias have occasionally occurred after intake of doses in excess of 200 mg.

Full prescribing information available on request