



If you do, you know why Nizoral *cream* is the right answer to **seborrhoeic dermatitis**

Apply twice daily until a few days after remission of symptoms.

Indications: Nizoral cream is indicated for topical application in the treatment of dermatophyte infections of the skin, as well as in the treatment of cutaneous candidosis and tinea (pityfissis) versicolor. Nizoral cream is also indicated for the treatment of sebornhoeic dermatilis, a skin condition related with the presence of Pityrosporum ovale. Precautions: Irritation may occur when Historia cream is used immediately after prolonged treatment with topical corticosteroids. Therefore, it is recommended to continue applying the topical corticosteroid in the mornings and to apply Nizoral cream in the evenings, and to subsequently and gradually withdraw the steroid therapy over a period of 2-3 weeks.



world leader in antimycotic research

PHARMACEUTICA B-2340 Beerse, Belgium "A man apart" . . . "A truly remarkable man" "

"Distinction of mind, of bearing, of character . . . characterised him always."

Ryle was an outstanding clinician and an original clinical investigator, his work on the fractional test meal and the introduction of "Ryle's tube" for taking gastric samples earning him an international reputation. With a holistic approach to medicine he saw the physician as a naturalist who must study how and why man reacts to disease as well as how disease progresses in man.

The awareness of a wider perspective led to his accepting the first chair of social medicine, at Oxford, where among his pupils was Michael Shepherd, now professor at the Institute of Psychiatry, who has written the introduction to the new Keynes Press edition of Ryle's The Natural History of Disease. First published in 1936, the collection of 34 lectures and essays is written in an unpretentiously clear and elegant style, setting out Ryle's observations on disease states and the principles underlying his work and outlook. Startingly relevant today, it is also, as Michael Shepherd says, a classic in its own right.

THE KEYNES PRESS



The Natural History of Disease

JOHN A RYLE

WITH AN INTRODUCTION BY MICHAEL SHEPHERD

THE KEYNES PRESS
BRITISH MEDICAL ASSOCIATION

The Natural History of Disease
consists of 400 numbered copies published by
the Keynes Press, London.
It has been designed by Sebastian Carter
and set in Garamond,
printed and bound by
Cambridge University Press
on a cream machine finished paper.

xxiv + 427 pages
Inland £50.00; Abroad £57.00; USA \$95.00
including postage.
Access, American Express, Visa credit cards accepted.
Please give full details.

Order from British Medical Journal (Keynes Press), PO Box 296, London WC1H 9TE.

Two new titles from the BMJ's series of books of general interest by medical writers





Portraits from Memory by Sir James Howie

One of the BMJ's most popular series, Sir James Howie's Portraits from Memory is now published as a complete collection. For this Memoir Club edition the author has provided three new portraits, an introductory self portrait, and an appendix in which he comments on the points raised in the widespread correspondence that resulted from the portraits' publication.

Sir James Howie's distinguished career has included appointments as professor of bacteriology in the University of Glasgow, director of the Public Health Laboratory Service, president of the Royal College of Pathologists,

and president of the British Medical Association. This book is not about himself, however, but about the memorable personalities he has known in microbiology and in military, hospital, and academic medicine. Essential reading for all students of human nature.


Price: Inland £14.95; Abroad £17.50; USA \$29.00. BMA members: Inland £13.95; Abroad £16.50; USA \$27.00.

A Natural History of Everyday Life: A biographical guide for would-be doctors of society by Lord Taylor of Harlow

As doctor, politician, and administrator the late Lord Taylor was a key figure in the formative years of some varied institutions: the National Health Service; Harlow New Town (playing an important part in its development and in its industrial health service); and the Memorial University of Newfoundland, of which he was the first president and vice-chancellor. He also laid the foundations of Medicare in Canada by his skilful negotiations to resolve the doctors' strike in Saskatchewan. In his lively and provocative memoirs Lord Taylor relates his "personal voyage" from childhood, through student days at St. Thomas's to his later career in public life, providing on the way innumerable pungent observations together with his views on the origins of some of society's ills—and their cures.

Price: Inland £19.95; Abroad £24.00; USA \$40.00


BMA members: Inland £18.95; Abroad £23.00; USA \$38.00.




ORDER FORM

BRITISH MEDICAL JOURNAL, PO Box 295, London WC1H 9TE or any leading medical bookseller.

		•
Please send me the following books:		□ I enclose
	□ Portraits from Memory	BMA Membership No.
The MEMOIR Club	☐ A Natural History of	Please make cheques payable to British Medical Journal or
	Everday Life	debit my credit card (please tick box)
	Prices include postage, by air abroad	□ Access □ American Express □ Mastercard □ Visa
		Card No.
Please print name and address clearly		Card expiry date
ADDRESS		• •
		Signature
•••••		□ Please send me a book catalogue.







Uses: Hismanal is a potent and non-sedative antihistamine indicated for the treatment of seasonal and perennial allergic rhinitis, allergic conjunctivitis, chronic urticaria and other allergic conditions.

Dosage and administration: adults and children over 12 years: 10 mg daily. Children 6-12 years: 5 mg once daily. Children younger than 6 years: 2 mg per 10 kg daily. Hismanal should be taken on an empty stomach. Precautions: Hismanal should be used in pregnant women only when, in the judgement of the physician, the potential benefits outweigh the possible hazards.

Adverse reactions: weight gain may occur during prolonged treatment. Overdose: in case of overdose, gastric lavage should be followed by close observation and ECG monitoring as arrhythmias have occasionally occurred after intake of doses in excess of 200 mg.