

TRADEMARK

(ketoconazole)

## Did you know

that in seborrhoeic dermatitis a dominant role is played by *Pityrosporum* yeasts?

## And did you know

that *Pityrosporum* is exceptionally sensitive to ketoconazole, alias Nizoral?

If you do, you know why Nizoral *cream* is the right answer to **seborrhoeic dermatitis**

Apply twice daily until a few days after remission of symptoms.

**Indications:** Nizoral *cream* is indicated for topical application in the treatment of dermatophyte infections of the skin, as well as in the treatment of cutaneous candidosis and tinea ( pityriasis) versicolor. Nizoral *cream* is also indicated for the treatment of seborrhoeic dermatitis, a skin condition related with the presence of *Pityrosporum ovale*. **Precautions:** Irritation may occur when Nizoral *cream* is used immediately after prolonged treatment with topical corticosteroids. Therefore, it is recommended to continue applying the topical corticosteroid in the mornings and to apply Nizoral *cream* in the evenings, and to subsequently and gradually withdraw the steroid therapy over a period of 2-3 weeks.



**JANSSEN**


*world leader in antimycotic research*

PHARMACEUTICA  
B-2340 Beerse, Belgium

CLINICAL EFFICACY PROVEN WORLDWIDE

# ROCEPHIN

FOR EXAMPLE: CLINICAL SUCCESS RATE  
IN MAJOR SERIOUS INFECTIONS



Reference

Data on file, F. Hoffmann-La Roche & Co. Limited Company, Basle, Switzerland.

Full details on composition, indications, contraindications, side effects, dosage and precautions are available on request.

ROCHE

F. HOFFMANN-LA ROCHE & CO. LIMITED COMPANY, BASLE, SWITZERLAND



Clearing  
his allergy  
without clouding his head

ONCE-A-DAY

TRADEMARK  
**Hismanal** astemizole  
The antihistamine of exceptional strength  
Effective from the first to the last day of therapy



**JANSSEN**  
PHARMACEUTICA

B-2340 Beerse, Belgium

**Prescribing Information**

**Uses:** Hismanal® is a potent and non-sedative antihistamine indicated for the treatment of seasonal and perennial allergic rhinitis, allergic conjunctivitis, chronic urticaria and other allergic conditions. **Dosage and administration:** adults and children over 12 years: 10 mg daily. Children 6-12 years: 5 mg once daily. Children younger than 6 years: 2 mg per 10 kg daily. Hismanal should be taken on an empty stomach. **Precautions:** Hismanal should be used in pregnant women only when, in the judgement of the physician, the potential benefits outweigh the possible hazards. **Adverse reactions:** weight gain may occur during prolonged treatment. **Overdose:** in case of overdose, gastric lavage should be followed by close observation and ECG monitoring as arrhythmias have occasionally occurred after intake of doses in excess of 200 mg.

Full prescribing information available on request