

TRADEMARK

(ketoconazole)

Did you know

that in seborrhoeic dermatitis
a dominant role is played by
Pityrosporum yeasts?

And did you know

that *Pityrosporum* is exceptional
sensitive to ketoconazole,
alias Nizoral?

Nizoral

If you do, you know why Nizoral cream is the right answer to seborrhoeic dermatitis

Apply twice daily until a few days after remission of symptoms.

Indications: Nizoral cream is indicated for topical application in the treatment of dermatophyte infections of the skin, as well as in the treatment of cutaneous candidosis and tinea (pityriasis) versicolor. Nizoral cream is also indicated for the treatment of seborrhoeic dermatitis, a skin condition related with the presence of *Pityrosporum ovale*. **Precautions:** Irritation may occur when Nizoral cream is used immediately after prolonged treatment with topical corticosteroids. Therefore, it is recommended to continue applying the topical corticosteroid in the mornings and to apply Nizoral cream in the evenings, and to subsequently and gradually withdraw the steroid therapy over a period of 2-3 weeks.



JANSSEN

PHARMACEUTICA

B-2340 Beerse, Belgium

world leader in antimycotic research

Full prescribing information
available on request.

IN LIFE-THREATENING INFECTIONS...

◁ROCEPHIN▷



...THE TREATMENT FOR EMERGENCIES

FOR EXAMPLE: SEPSIS... PATHOGENS ERADICATED IN 96.3% OF ALL CASES

^{Reference}

^{*}Data on file, F. Hoffmann-La Roche & Co. Limited Company, Basle, Switzerland.

Full details on composition, indications, contraindications, side effects, dosage and precautions are available on request.



F. HOFFMANN-LA ROCHE & CO. LIMITED COMPANY, BASEL, SWITZERLAND

Adalat® The cardioprotective coronary therapeutic agent



Adalat® Nifedipine: capsules 10 mg. **Indications:** CHD: chronic stable angina pectoris (stress-induced angina), unstable, including vasospastic angina, post-infarction angina (start treatment 8 days after acute episode). **Hypertension. Dosage:** According to severity of disease and indication. Generally 1 capsule 3 x daily; could be increased to 2 capsules 3 x daily. For coronary spasms temporarily up to 2 capsules 4 – 6 x daily. If rapid onset required: bite through and swallow 1 – 2 capsules; see full prescribing information. Single dosage interval of 2 capsules never less than 2 hours. **Contraindications:** Hypersensitivity to active substance; pregnancy; lactation; cardiovascular shock. **Precautions:** Severe hypotension; dialysis patients with malignant hypertension: close monitoring. **Interactions:** Antihypertensive agents, cimetidine: may enhance the antihypertensive effect. Same with beta-blockers; possible development of heart failure. **Side effects:** Often mild and transient vasodilation, hypotensive, skin and other reactions; see full prescribing information. Solitary gingival hyperplasia, chest pain; if relation to Adalat established, discontinue therapy. Extremely rare liver function disturbances (including intrahepatic cholestasis), temporary hyperglycaemia, gynaecomastia (regression on drug discontinuation). Ability to drive or operate machinery may be impaired.

Full prescribing information available from Bayer AG, Leverkusen, West Germany.

Co.-No.: 2

