Full prescribing information available on request.

Did you know that in seborrhoeic dermatitis a dominant role is played by Pityrosporum yeast\$? And did you kn that *Pityrosporum* is exceptional sensitive to ketoconazole, alias Nizoral?

If you do, you know why Nizoral *cream* is the right answer to **seborrhoeic dermatitis**

Apply twice daily until a few days after remission of symptoms.


Indications: Nizoral crasm is indicated for topical application in the treatment of dermatophyte infections of the skin, as well as in the treatment of cutaneous candidosis and tinea (pityriasis) versicolor. Nizoral cream is also indicated for the treatment of seborthoeic dermatkis, a skin condition related with the presence of Pityrosporum ovale. Precautions: Irritation may occur when Nizoral cream is used immediately after prolonged treatment with topical corticosteroids. Therefore, it is recommended to continue applying the topical corticosteroid in the mornings and to apply Nizoral cream in the evenings, and to subsequently and gradually withdraw the stand therapy over a period of 2-3 weeks.



JANSSER

world leader in antimycofic research

PHARMACEUTICA B-2340 Beerse, Belgium





HISManal The antihistamin

Effective from the first to the last day of therapy



Prescribing information

Uses: Hismanal' is a potent and non-sedative antihistamine indicated for the treatment of seasonal and perennial allergic conjunctivitis, chronic urticaria and other allergic conditions.

Dosage and administration: adults and children over 12 years: 10 mg daily. Children 6-12 years: 5 mg once daily. Children younger than 6 years: 2 mg per 10 kg daily. Hismanal should be taken on an empty stomach. Precautions: Hismanal should be used in pregnant women only when, in the judgement of the physician, the potential benefits outweigh the possible hazards.

Adverse reactions: weight gain may occur during prolonged treatment. Overdose: in case of overdose, gastric lavage should be followed by close observation and ECG monitoring as arrhythmias have occasionally occurred after intake of doses in excess of 200 mg.