Some patients are more susceptible to vaginal candidosis than others



Candida cells are capable of penetrating to the depth of several layers of the vaginal epithelium. This suggests that these hidden yeasts may be protected from topical antifungal agents, only to re-emerge and proliferate again some time later when the epithelial cells are normally shed. The deeper layers of the vaginal mucosa are more accessible by systemic than by topical route. Thus, in patients who appear highly susceptible to vaginal candidosis. oral Nizoral treatment makes good sense.

NIZOTA (ketoconazole

2 oral tablets once daily for 5 days

Full prescribing information available on request.

Prescribing notes: For maximal absorption Nizoral should be taken with meals. Nizoral is contra-indicated in pregnancy. Precautions: the use of agents which reduce gastric acidity (anti-cholinergic drugs, antacids, H₂-blockers) should be avoided and, if indicated, such drugs should be taken not less than 2 hours after Nizoral. Side-effects: nausea, skin rash, headache and pruritus may occasionally be observed. Alterations in liver function tests have occurred in patients on Nizoral; these changes may be transient. Cases of hepatitis have been reported with an incidence of about 1 per 10,000 patients. If a patient develops jaundice or any symptoms suggestive of hepatitis, treatment with Nizoral should be stopped. Mild asymptomatic increases of liver enzyme levels, on the other hand, do not necessitate discontinuation of the treatment.



world leader in antimycotic research



Price: Inland £7.95; Abroad £9.50 BMA members: Inland £7.45; Abroad: £9.00

Just published

Many medical journals, including the British Medical Journal, now expect scientific papers submitted to them to contain confidence intervals when appropriate. Why? what are they? and how do you calculate them? Statistics with Confidence tells you. A clear explanation of the reasons for using confidence intervals is followed by detailed presentation of methods of calculation, including numerous worked examples and specially compiled tables. To make things even easier, a computer programme, Confidence Interval Analysis (CIA), for calculating confidence intervals, has been specially designed by Martin Gardner and details are available from the Publishing Department, British Medical Journal.

Eighth edition

The statistical testing of data is indispensable in many types of medical investigation and a help on countless occasions in clinical practice. This book provides step by step instruction. Subjects covered include standard deviation, χ^2 tests, t tests, non-parametric tests, and correlation. The book includes sections on Fisher's exact probability test and rank correlation not published in the original BMJ series. Methods specially adapted to pocket calculators.



Price: Inland £3.00; Abroad £4.50 BMA members: Inland £2.50; Abroad: £4.00

"He uses statistics as a drunken man uses lamp-posts for support rather than illumination." *

Let the BMF throw light on your methods and help your results stand up

* Andrew Lang

STATISTICS IN PRACTICE STATISTICS IN STATISTICS AND ETHICS IN MEDICAL RESEARCH Douglas of American Medical American Company of American Company o


Price: Inland £8.95; Abroad £11.00 BMA members: Inland £7.45; Abroad: £10.00

Fourth impression

No doctor can afford to ignore statistics: most modern medical research uses statistics. This important and authoritative book provides clear information on designing studies, applying statistical techniques, and interpreting studies that use statistics. It can be easily understood by those with no statistical training and should be read by all those who want to keep abreast of new developments.

Second edition

Epidemiology has its own techniques of data collection and interpretation and its necessary jargon of technical terms, and in Epidemiology for the Uninitiated Professors Geoffrey Rose and David Barker guide the novice expertly through the theory and practical pitfalls. The second edition of this popular BMJ handbook has been revised to include further details of epidemiological methods and some of their more dramatic applications, such as the investigations on the Spanish cooking oil epidemic, and AIDS.





Price: Inland £3.95; Abroad £5.50 BMA members: Inland £3.45; Abroad: £5.00

ORDER FORM

BRITISH MEDICAL JOURNAL, P.O. Box 295, London WC1H 9TE, England or any leading medical bookseller

Please send me	I enclose BMA Membership No
 □ Statistics with Confidence □ Statistics at Square One □ Statistics in Practice 	Please make cheques payable to British Medical Journal or debit my credit card (please tick box)
☐ Epidemiology for the Uninitiated All prices include postage, by air abroad	☐ Access/Mastercard ☐ Barclaycard/Visa ☐ American Express
NAME	Card No.
ADDRESS	CARD EXPIRY DATE
	SIGNATURE
POSTCODE	☐ Please send me a book catalogue







ONCE-A-DAY ismana

The antihistamine of exceptional strength Effective from the first to the last day of therapy



Uses: Hismanal* is a potent and non-sedative antihistamine indicated for the treatment of seasonal and perennial allergic conjunctivitis, chronic urticaria and other allergic conditions.

Dosage and administration: adults and children over 12 years: 10 mg daily. Children 6-12 years: 5 mg once daily. Children younger than 6 years: 2 mg per 10 kg daily. Hismanal should be taken on an empty stomach. Precautions: Hismanal should be used in pregnant women only when, in the judgement of the physician, the potential benefits outweigh the possible hazards.

Adverse reactions: weight gain may occur during prolonged treatment. Overdose: in case of overdose, gastric lavage should be followed by close observation and ECG monitoring as arrhythmias have occasionally occurred after intake of doses in excess of 200 mg.