Some patients are more susceptible to vaginal candidosis than others

Candida cells are capable of penetrating to the depth of several layers of the vaginal epithelium. This suggests that these hidden yeasts may be protected from topical antifungal agents, only to re-emerge and proliferate again some time later when the epithelial cells are normally shed. The deeper layers of the vaginal mucosa are more accessible by systemic than by topical route. Thus, in patients who appear highly susceptible to vaginal candidosis, oral Nizoral treatment makes good sense.

Nizoral (Lotocopa)

2 oral tablets
once daily for 5 days


Full prescribing information available on request.

Prescribing notes: For maximal absorption Nizoral should be taken with meals. Nizoral is contra-indicated in pregnancy. Precautions: the use of agents which reduce gastric acidity (anti-cholinergic drugs, antacids, $\rm H_2$ -blockers) should be avoided and, if indicated, such drugs should be taken not less than 2 hours after Nizoral. Side-effects: nausea, skin rash, headache and pruritus may occasionally be observed. Alterations in liver function tests have occurred in patients on Nizoral; these changes may be transient. Cases of hepatitis have been reported with an incidence of about 1 per 10,000 patients. If a patient develops jaundice or any symptoms suggestive of hepatitis, treatment with Nizoral should be stopped. Mild asymptomatic increases of liver enzyme levels, on the other hand, do not necessitate discontinuation of the treatment.

JANSSEN PHARMACEUTICA
B-2340 Beerse, Belgium

world leader in antimycotic research

J.PH. 198



Essays to honour Ian Munro

Is the *Lancet* the most famous medical journal in the world?

It is certainly the most frequently cited, its mixture of high science, common sense, mild subversion, and humour giving it a loyal following throughout the world. Much of this is due to the character of its editor, Ian Munro.

Taken on to the editorial staff because his epistolary style impressed the then editor, Sir Theodore Fox, Munro became the longest serving medical staff member of a medical journal.

He has recently retired, and to mark the occasion friends and colleagues at home and abroad have produced a

literary substitute for the traditional carriage clock. In it they pay tribute to Munro's qualities as editor, cricketer, liberal campaigner, and soother of hurt authors' feelings.

Contributors include the editors of the Norwegian Medical Journal, the New Zealand Medical Journal, and the BMJ as well as the deputy editors of the Lancet. You don't have to be Ian Munro to read and enjoy this book: a limited edition of 100 copies is now published by the Keynes Press.

Designed and printed by Sebastian Carter at the Rampart Lions Press, Cambridge, on Zerkall laid mould made paper. The text is set in Monotype Perpetua by Monoset Ltd. and the

The text is set in Monotype Perpetua by Monoset Ltd, and the displayed matter in Golden Cockerel roman, both designed by Eric Gill. Thread sewn in a soft cover.

48 pages + 5 pages of plates.

Price:

Inland £20.00; Abroad £25.00, including postage

Access, American Express and Visa credit cards accepted. Please give full details.







ORDER FROM:

British Medical Journal (The Keynes Press), PO Box 295, London WC1H 9TE.

Just published

High tech equipment for diagnosis and treatment is rightly valued, but for the many disabled people in the population it is often relatively simple aids that make all the difference to the quality of life. Do you know what they are? Where they can be obtained? Which ones are suitable and effective? In Everyday Aids and Appliances the authors give critical evaluation of some of the various types of aids available: those attached to the body (such as hearing aids, stomas, special footwear); those held but not necessarily fixed to the body (such as mobility aids); and equipment in the home which allows independence in everyday activities (such as wheelchairs, hoists).





PO Box 295,

BRITISH MEDICAL JOURNAL

(a) Heel wedge; (b) elongated heel; (c) rocker sole; (d) floated out heel.

ORDER FORM

Please send me	London WC1H 9TE, or any leading bookseller
Flease send me ☐ Everyday Aids and	Annliances
and the following books:	
NAME	
l	
ł	
1	

I enclose BMA Membership No.
Please make cheques payable to British Medical Journal

☐ Access ☐ Visa ☐ American Express ☐ Mastercard

or debit my credit card (please tick box)

☐ Please send me a book catalogue

Card No.

Card expiry date

Originally published in the British Medical Journal and now collected in one volume with additional illustrations, these articles bring you fully up to date on items that may greatly benefit disabled people and those who care for them.



Price: Inland £5.95; Abroad £7.00

BMA Members: Inland £5.45;

Abroad £6.50

including postage, by air abroad



The abstract theory of theories has highly practical applications, and the authors explore the uses of computer technology and artificial intelligence systems as well as "fuzzy logic," "relevant logic," and the logic of economics and ethics. A book for all who wish to clear their minds of cant.

Price: Inland £5.95; Abroad £7.50 BMA members: Inland £5.45; Abroad £7.00 (including postage, by air abroad)


Payment must be enclosed with order


ORDER YOUR COPY NOW FROM British Medical Journal, PO Box 295 London WC1H 9TE or any leading medical bookseller


Think about it

You make a clinical diagnosis. Do you ever consider the thought processes by which you arrived at it? Medical students and practitioners are often concerned with examples of diagnostic logic, but seldom consider them in the context of a general philosophy. Is diagnostic logic out on a limb, or is it based on the same principles as logic in general?

In Logic in Medicine doctors and philosophers combine to provide a coherent system of diagnostic logic with a broader view of the science and art of reasoning.








Adalat®retard:

The antihypertensive from Bayer with cardioprotective effect. Worldwide.



crisis, Adalat <u>capsules</u> (10 mg) should be used; see full prescribing information. <u>Dosage</u>: Generally 1 tablet twice daily; in some cases 2 tablets twice daily; see full prescribing information. Single dosage interval of 2 tablets never less than 4 hours. <u>Contraindications</u>: Hypersensitivity to active substance; pregnancy; lactation; cardiovascular shock. <u>Precautions</u>: Severe hypotension; dialysis patients with malignant hypertension: close monitoring. <u>Interactions</u>: Antihypertensive agents, cimetidine: may enhance the antihypertensive effect. Same with beta-blockers: possible development of heart failure. Side effects: Often mild and transient vasodilation, hypotensive,

skin and other reactions; see full prescribing information. Solitary gingival hyperplasia, chest pain; if relation to Adalat established, discontinue therapy. Extremely rare liver function disturbances (including intrahepatic cholestasis), temporary hyperglycaemia, gynaecomastia (regression on drug discontinuation). Ability to drive or operate machinery may be impaired.

Full prescribing information available from Bayer AG, Leverkusen, West Germany.

bayer