Some patients are more susceptible to vaginal candidosis than others



Candida cells are capable of penetrating to the depth of several layers of the vaginal epithelium.

This suggests that these hidden yeasts may be protected from topical antifungal agents, only to re-emerge and proliferate again some time later when the epithelial cells are normally shed.

The deeper layers of the vaginal mucosa are more

accessible by systemic than by topical route. Thus, in patients who appear highly susceptible to vaginal candidosis, oral Nizoral treatment makes good sense.

NZOTA
(ketoconażole

2 oral tablets once daily for 5 day


Full prescribing information available on request.

Prescribing notes: For maximal absorption Nizoral should be taken with meals. Nizoral is contra-indicated in pregnancy. Precautions: the use of agents which reduce gastric acidity (anti-cholinergic drugs, antacids, H₂-blockers) should be avoided and, if indicated, such drugs should be taken not less than 2 hours after Nizoral. Side-effects: nausea, skin rash, headache and pruritus may occasionally be observed. Alterations in liver function tests have occurred in patients on Nizoral; these changes may be transient. Cases of hepatitis have been reported with an incidence of about 1 per 10,000 patients. If a patient develops jaundice or any symptoms suggestive of hepatitis, treatment with Nizoral should be stopped.


Mild asymptomatic increases of liver enzyme levels, on the other hand, do not necessitate discontinuation of the treatment.


JANSSEN PHARMACEUTICA
B-2340 Beerse, Belgium

world leader in antimycotic research Genetics is a subject fundamental to medicine, but rapid advances in recent years may make the subject complex and confusing to clinicians and paramedical staff outside the specialty. Nevertheless, the demand for genetic investigation and counselling is increasing, and these services need to be offered by clinicians other than clinical geneticists. In the ABC of Clinical Genetics Helen Kingston, consultant clinical geneticist at St. Mary's Hospital, Manchester, reviews basic concepts of inheritance in mendelian, chromosomal, and multifactorial disorders; outlines the investigation and management of genetic conditions; and illustrates the application of recombinant DNA technology to clinical practice. Implications of genetic diseases for relatives are also considered, together with the ethical and human dilemmas that face doctors and patients and their families.




ORDER FORM	BRITISH MEDICAL JOURNAL PO Box 295, London WC1H 9TE, or any leading bookseller	
Please send me		
☐ ABC OF CLINICAL GENETICS		
and the following BMJ books:		
1		
	•••••	
NAME		
ADDRESS		
l		
POSTCODE		
I enclose	BMA Membership No	
Please make cheques pa or debit my credit card	yable to British Medical Journal (please tick box)	
│ □ Visa □ American	Express	
Card No.		
Card expiry date		
Signature		
☐ Please send me a boo	ok catalogue	





Inland £10.95; Abroad £13.00 BMA Members: Inland £9.95; Abroad £12.00 including postage, by air abroad

You don't know everything





But often in a medical career it seems as if you are expected to. When confronted with a problem you've never met before the only thing to do is to ask the experts—but where are the experts who are never on holiday, never off sick, in a meeting, or simply unavailable? In *How To Do It: 3*, that's where. You can keep them in your pocket, and in a few minutes they can tell you, among many other things, how to:

- lecture overseas
- be a manager
- prepare a Festschrift
- commission a portrait
- set up an audit
- write a practice annual report
- organise multicentre trials
- start a DNA diagnostic service
- survive a dinner

Two companion volumes of essential know-how are also available


Price (each book): Inland £6.95; Abroad £9.00 BMA members: Inland £6.45; Abroad £8.50





ORDER FORM British Medical Journal, PO Box 295, London WC1H 9TE, England or any leading bookseller		
Please send me	copy/copies of	☐ I enclose BMA Membership No
 ☐ HOW TO DO IT: 2 ☐ HOW TO DO IT: 2 ☐ HOW TO DO IT: 1 	2	Please make cheques payable to British Medical Journal or debit my credit card (please tick box)
Prices to include postage, by		\square visa \square american express \square mastercard
NAME		CARD NUMBER
. ,		Card expiry date
		Signature
POST CODE		☐ Please send me a book catalogue

The Royal Society of Medicine




There are many reasons for joining The Royal Society of Medicine – here are six of them:

- * The Royal Society of Medicine has 34 specialty sections holding 250 high-level meetings annually at home and abroad...
- * The Royal Society of Medicine's library is one of the largest medical collections in Europe...
- * The Royal Society of Medicine publishes a monthly Journal sent free to all members...
- * The Royal Society of Medicine's club facilities, including restaurants and bar, are in a modern, elegant setting...
- * The Royal Society of Medicine provides hotel-standard accommodation for members at *less than half* London hotel prices...
- * The Royal Society of Medicine has its own television channel...

New members are welcome

For further details, please write to:
Jill Thomas
Information Secretary (3)
The Royal Society of Medicine
Wimpole Street, London W1M 8AE
or ring 01-408 2119, Ext. 279





Many medical journals, including the British Medical Journal now expect scientific papers submitted to them to contain confidence intervals when appropriate. Why? what are they? and how do you calculate them? Statistics with Confidence tells you. A clear explanation of the reasons for using confidence intervals is followed by detailed presentation of methods of calculation, including numerous worked examples and specially compiled tables. To make things even easier, a computer programme, Confidence Interval Analysis (CIA), for calculating confidence intervals has been specially designed by Martin Gardner and details are available from the Publishing Department, British Medical Journal.

An essential handbook for everybody using statistical methods to present their findings (and that means most doctors). Statistics with Confidence can also be applied to other disciplines—such as engineering, agriculture, and the social sciences.

Price: Inland £7.95; Abroad £9.50

BMA Members: Inland £7.45; Abroad £9.00

including postage, by air abroad

Order from: BRITISH MEDICAL JOURNAL, PO BOX 295, LONDON WC1H 9TE or any leading bookseller. Please enclose payment with order, or send us full details of your ACCESS, VISA or AMERICAN EXPRESS credit card.

Render to

histamine

what histamine deserves

Histamine deserves the antihistamine that can really take it on. For right away the allergies that thrive on histamine will respond.

Typical examples are hay fever, perennial rhinitis, allergic conjunctivitis and different forms of urticaria.

Hismanal is unique by the strength of its histamine-antagonism.

Without any risk of sedation, its full antihistamine strength can be utilized.

And there is no weakening during the 24 hour dosing interval, so that the symptoms won't get a chance to cause untimely trouble.

Hismanal

The antihistamine of exceptional strength.

Effective from the first to the last day of therapy.





the drug discovery company

Prescribing information

Uses: Hismanal is a potent and non-sedative antihistamine indicated for the treatment of seasonal and perennial allergic rhinitis, allergic conjunctivitis, chronic urticaria and other allergic conditions.

Dosage and administration: adults and children over 12 years: 10 mg daily. Children 6-12 years: 5 mg once daily. Children younger than 6 years: 2 mg per 10 kg daily. Hismanal should be taken on an empty storrach. Precautions: Hismanal should be used in pregnant women only when, in the judgement of the physician, the possible hazards.

Adverse reactions: weight gain may occur during prolonged treatment. Overdose: in case of overdose, gastric lavage should be followed by close observation and ECG monitoring as antitythmias have occasionally occurred after intake of doses in excess of 200 mg.

uil prescribing information available on request.



CHOST AND CONTRACT

898