

TRADEMARK
Sporanox
ITRACONAZOLE

THE VERSATILE ORAL ANTIFUNGAL

Sporanox is highly active against virtually all dermatophytes and yeasts, and against notoriously virulent pathogenic fungi, such as *Aspergillus* spp., some *Fusarium* spp. and *Sporothrix schenckii*.

That's why, in dermatology, gynaecology and internal medicine alike, Sporanox is rapidly becoming the oral treatment of choice for troublesome fungal infections.

Simplicity itself:

- in tinea infections of the skin:
1 capsule daily for precisely 15 days*
- in vaginal candidosis:
2 capsules b.i.d. for only 1 day

* Involvement of highly keratinized skin such as hand palms and foot soles requires an additional 15-day treatment.

Note: Sporanox (itraconazole) is not yet available in all countries.

Administration: It is essential that Sporanox be taken immediately after a full meal for maximal absorption. **Contra-indications:** Sporanox is contra-indicated during pregnancy. Adequate contraceptive precautions should be taken by women of childbearing potential during the menstrual cycle of Sporanox therapy. **Warnings and precautions:** Sporanox is predominantly metabolized in the liver. Although clinically Sporanox has not been associated with hepatic dysfunction, it is advisable not to give this drug to patients with a known history of liver disease or to patients who have experienced liver toxicity with other drugs. Nursing mothers: it is recommended not to breast feed whilst taking Sporanox. Paediatric use: Sporanox has not been systematically studied in children; it should, therefore, not be used in paediatric patients unless the potential benefit outweighs the potential risks. **Drug interactions:** Sporanox should not be given concomitantly with rifampicin. **Adverse reactions:** Side-effects during Sporanox therapy occurred in 7.1% of the patients. The most frequently reported side-effects were nausea (1.3%), abdominal pain (1.2%), headache (1%) and dyspepsia (0.7%). **Supplied:** Blister packs of 4, 6, or 15 capsules. Each capsule contains 100 mg of itraconazole.

Full prescribing information is available on request.

Trichophyton rubrum


Candida albicans
Sabouraud agar

Aspergillus fumigatus

Fusarium solani

Sporothrix schenckii

Cryptococcus neoformans




Abbott
Laboratories


CHILDREN ARE SENSITIVE

«ROCEPHIN»

**PATHOGENS
ELIMINATED**



**No
SIDE EFFECTS**



WELL TOLERATED EVEN IN CHILDREN
(Analysis of literature on 887 patients)

Components: Ceftriaxone. **Indications:** Sepsis; meningitis; abdominal infections; infections of the bones/joints, soft tissue, skin and of wounds; infections in patients with impaired defence mechanisms; renal and urinary tract infections; respiratory tract infections, particularly pneumonia, and ear, nose and throat infections; genital infections, including gonorrhoea. Perioperative prophylaxis of infections. **Dosage:** Usual dosage for adults: 1-2 g once daily (every 24 hours). Further information on administration and special dosage

recommendations are available on request. **Contra-indications:** Known hypersensitivity to cephalosporins. **Precautions:** Pregnancy (particularly in the first trimester) unless absolutely necessary. Hypersensitivity to penicillin (possibility of allergic cross-reactions). Anaphylactic shock. **Side effects:** Gastrointestinal complaints. Hematological changes. Skin reactions. **Packs:** Packs for i.m. injection, 5's. Packs for i.v. injection, 5's. Packs for i.v. infusion, 5's. Full details are available on request.


Reference: Data on file, F. Hoffmann-La Roche Ltd, Basel, Switzerland



F. Hoffmann-La Roche Ltd, Basel, Switzerland

THE ANTIHISTAMINE of exceptional strength

TRADEMARK
Hismanal
(astemizole)



Prescribing information: Hismanal is a potent antihistamine indicated for the treatment of seasonal and perennial allergic rhinitis, allergic conjunctivitis, and other allergic conditions. **Dosage and administration:** adults, 5 mg once daily. Children younger than 6 years, 1 mg once daily. **Precautions:** Hismanal should be used in patients with liver disease. If the potential benefits outweigh the possible hazards, the physician should monitor the patient during prolonged treatment. **Full prescribing information**

JUST
ONE TABLET
A DAY



company