

TRADEMARK

Sporanox

ITRACONAZOLE


THE VERSATILE ORAL ANTIFUNGAL

Sporanox is highly active against virtually all dermatophytes and yeasts, and against notoriously virulent pathogenic fungi, such as *Aspergillus* spp., some *Fusarium* spp. and *Sporothrix schenckii*.

That's why, in dermatology, gynaecology and internal medicine alike, Sporanox is rapidly becoming the oral treatment of choice for troublesome fungal infections.

Sporanox

4 gelules capsules



Simplicity itself:

- in tinea infections of the skin:
1 capsule daily for precisely 15 days*
- in vaginal candidosis:
2 capsules b.i.d. for only 1 day

* Involvement of highly keratinized skin such as hand palms and foot soles requires an additional 15-day treatment.

Note: SporanoX (itraconazole) is not yet available in all countries.

Administration: It is essential that SporanoX be taken immediately after a full meal for maximal absorption. **Contra-indications:** SporanoX is contra-indicated during pregnancy. Adequate contraceptive precautions should be taken by women of childbearing potential during the menstrual cycle of SporanoX therapy. **Warnings and precautions:** SporanoX is predominantly metabolized in the liver. Although clinically SporanoX has not been associated with hepatic dysfunction, it is advisable not to give this drug to patients with a known history of liver disease or to patients who have experienced liver toxicity with other drugs. Nursing mothers: it is recommended not to breast feed whilst taking SporanoX. Paediatric use: SporanoX has not been systematically studied in children; it should, therefore, not be used in paediatric patients unless the potential benefit outweighs the potential risks. **Drug interactions:** SporanoX should not be given concomitantly with rifampicin. **Adverse reactions:** Side-effects during SporanoX therapy occurred in 7.1% of the patients. The most frequently reported side-effects were nausea (1.3%), abdominal pain (1.2%), headache (1%) and dyspepsia (0.7%). **Supplied:** Blister packs of 4, 6, or 15 capsules. Each capsule contains 100 mg of itraconazole.

Full prescribing information is available on request.

*Trichophyton
rubrum*

Candida albicans
Sabouraud agar

*Aspergillus
fumigatus*

Fusarium solani

*Sporothrix
schenckii*

*Cryptococcus
neoformans*




Janssen Pharmaceutica
Beerse, Belgium
Center for Antimicrobial Research

CHILDREN ARE SENSITIVE


«ROCEPHIN»

24-HOUR BACTERICIDAL POWER

**PATHOGENS
ELIMINATED**



**No
SIDE EFFECTS**



WELL TOLERATED EVEN IN CHILDREN
(Analysis of literature on 887 patients)

Components: Ceftriaxone. **Indications:** Sepsis; meningitis; abdominal infections; infections of the bones, joints, soft tissue, skin and of wounds; infections in patients with impaired defence mechanisms; renal and urinary tract infections; respiratory tract infections, particularly pneumonia, and ear, nose and throat infections; genital infections, including gonorrhea. Perioperative prophylaxis of infections. **Dosage:** Usual dosage for adults: 1-2 g once daily (every 24 hours). Further informations on administration and special dosage

recommendations are available on request. **Contra-indications:** Known hypersensitivity to cephalosporins. **Precautions:** Pregnancy (particularly in the first trimester) unless absolutely necessary. Hypersensitivity to penicillin (possibility of allergic cross-reactions). Anaphylactic shock. **Side effects:** Gastrointestinal complaints. Hematological changes. Skin reactions. **Packs:** Packs for i.m. injection, 5's. Packs for i.v. injection, 5's. Packs for i.v. infusion, 5's. Full details are available on request.

Reference: Data on file, F. Hoffmann-La Roche Ltd, Basel, Switzerland



F. Hoffmann-La Roche Ltd, Basel, Switzerland

Render to

histamine

what histamine deserves



Histamine deserves the antihistamine that can really take it on. For right away the allergies that thrive on histamine will respond. Typical examples are hay fever, perennial rhinitis, allergic conjunctivitis and different forms of urticaria.

Hismanal is unique by the strength of its histamine-antagonism. Without any risk of sedation, its full antihistamine strength can be utilized.

And there is no weakening during the 24 hour dosing interval, so that the symptoms won't get a chance to cause untimely trouble.



Hismanal[®]

(astemizole)

The antihistamine
of exceptional strength.

Effective from
the first
to the last day
of therapy.



JANSSEN
PHARMACEUTICA

the drug discovery company
B-2340 Beerse, Belgium

Prescribing information

Uses: Hismanal is a potent and non-sedative antihistamine indicated for the treatment of seasonal and perennial allergic rhinitis, allergic conjunctivitis, chronic urticaria and other allergic conditions.

Dosage and administration: adults and children over 12 years: 10 mg daily. Children 6-12 years: 5 mg once daily. Children younger than 6 years: 2 mg per 10 kg daily. Hismanal should be taken on an empty stomach. **Precautions:** Hismanal should be used in pregnant women only when, in the judgement of the physician, the potential benefits outweigh the possible hazards.

Adverse reactions: weight gain may occur during prolonged treatment. **Overdose:** in case of overdose, gastric lavage should be followed by close observation and ECG monitoring as arrhythmias have occasionally occurred after intake of doses in excess of 200 mg.

Full prescribing information available on request.

JUST
ONE TABLET
A DAY