KEY WORDS OF MODERN ANTIFUNGAL THERAPY

THE ADVANTAGE LIPOPHILIC OF BEING



Dermatophytosis caused by Trichophyton concentricum

Fungi have a strong affinity for the lipid-rich layers of the skin, the mucosa and other tissues. In addition, fungal cell membranes consist largely of lipids.

Since most antifungal drugs are targeted at the fungal cell membranes, it is advantageous if they too are lipophilic in nature. Its lipophilicity is what helps an oral drug like itraconazole to exert its antifungal effect precisely where it is needed: in the fungal membranes and in the target tissues.

It also helps that itraconazole is decidedly keratinophilic. For this is why it is strongly attracted to the skin's stratum corneum where many fungi find the keratin they need to subsist.

Possessing both properties gives itraconazole the additional advantage that it remains in the epithelial cells for as long as it takes these cells to be desquamated. Its antifungal activity will therefore continue for several days or even weeks after stopping treatment, thus permitting oral dosage schedules to be limited to a short period of time.

In other words, in much the same way as we have become accustomed to using oral antibiotics, we can now also combat fungal infections with **short**, **fixed oral treatment schedules**.



SHORT AND SIMPLE ORAL THERAPY

(See prescribing information below)

Basic dose in dermatomycoses: 1 capsule (100 mg) once daily for 15 days Standard dose in vaginal candidosis: 2 x 2 capsules (400 mg) for 1 day only

* Trademarks: SPORANOX, SEMPERA, TRISPORAL

PHARMACEUTICA
B-2340 Beerse, Belgium
expertise in
antimycotic research


JANSSEN PHARMACEUTICA 1991

Properties: Sporanox (ftraconazole), a triazole derivative, is orally active against infections with dermatophytes (*Trichophyton spp., Microsporum spp., Epidermophyton floccosum*), yeasts (*Candida spp., Pityrosporum spp.*), Aspergiillus spp. and various other yeasts and trungi. Indications: Sporanox (ftraconazole) is indicated for vulvovaginal candidosis, pityriasis versicolor, dermatophytoses, fungal keratitis and oral candidosis. **Dosage and administration:** Vulvovaginal candidosis: 2 capsules (200 mg) morning and evening for 1 day; pityriasis


versicolor: 2 capsules (200 mg) once daily for 7 days; tinea corporis, tinea cruris, tinea pedis, tinea manus: 1 capsule (100 mg) daily for 15 days; highly keratinized regions, as in plantar tinea pedis and palmar tinea manus, require 1 capsule (100 mg) daily for 30 days. Oral candidosis: 1 capsule (100 mg) daily for 30 days. Fungal keratitis: 2 capsules (200 mg) once daily for 21 days. Contra-indications: Sporanox (traconazole) is contra-indicated during pregnancy. Warnings and precautions: Although clinically Sporanox (traconazole) has

not been associated with hepatic dysfunction, it is advisable not to give this drug to patients with a known history of liver disease. **Nursing mothers:** It is recommended not to breast feed whilst taking Sporanox (traconazole). **Drug interactions:** Sporanox (traconazole) should not be given concomitantly with rifampicin.

Full prescribing information is available on request.



Search 25 Years of the BMJ in Seconds



Save time and money

with Maxwell's new BMJ CD-ROM. It gives you thousands of BMJ

references, abstracts and articles all on a single disc. It's the most complete, easy-to-use research tool of its kind.

- More comprehensive 25 years of BMJ MEDLINE references and abstracts plus five years of the most recent articles including tables. Updated annually
- More search options Search by keyword or phrase, author and/or title using powerful BRS search software
- Easier to use Find a MEDLINE citation and immediately link to the article, plain-language prompts, tutorial, free technical support
- Available now All paid orders shipped within 48 hours

Operates on standard IBM PC 286 or 386 and compatibles with a CD-ROM drive. Call for special savings on CD-ROM hardware.


Build a compact library with other medical CD-ROMs from Maxwell Electronic Publishing: *The Lancet, New England Journal of Medicine, Annals of Internal Medicine, Morbidity and Mortality Weekly Reports, Journal of the American Medical Association*, AIDS, Viral Hepatitis.

	ROM
Institutional use US	\$440.
7 VISA □ MC □ .	AMEX
Exp. Date	
)M hardware.	pact Libraries
<u> </u>	
<u> </u>	
MAXWE	A0601
	RTER OF UBSCRIPTI Vell's BMJ CD- rice. Institutional use US regularly US\$550) VISA MC MC

For immediate delivery, Fax: 010-617-868-7738

Phone: 010-617-661-2955

gastric distress & oesophagitis hyperacidity or dysmotility?



restores upper G.I. motility like no other agent.

JANSSEN PHARMACEUTICA B-2340 Beerse, Belgium