

FUNGICIDE FROM **Sporanox®** itraconazole 100 mg

One of the notorious problems with fungal infections of the skin or the vagina is that the organism may penetrate the deeper layers of the epithelium, out of reach of topical medication. And besides, when treating fungal skin lesions locally, the infection is often already subclinically present at other sites of the body, waiting for a chance to start the trouble all over again.

Because Sporanox works orally, i.e. "from the inside out", it will destroy even the best hidden fungal cells. All the more so, because Sporanox has a strong affinity for epidermal and mucosal tissues as well as for the fungal cell wall itself where it must exert its fungicidal activity.

SHORT AND SIMPLE ORAL THERAPY

Standard dose in Dermatology: 1 capsule (100 mg) once daily for 15 days (Sporanox will remain active in the stratum corneum for another 3-4 weeks)

Standard dose in Gynaecology: 2 x 2 capsules (400 mg) for 1 day only
(Sporanox will remain active in the vaginal epithelium for another 3 to 4 days)

This product is not yet available in all countries.

* Trademarks: SPORANOX, SEMPERA,
TRISPORAL, SPORAL.

Properties: Sporanox (itraconazole), a triazole derivative, is orally active against infections with dermatophytes (*Trichophyton* spp., *Microsporum* spp., *Epidermophyton* spp.) and yeasts (*Candida* spp., *Aspergillus* spp., *Phycomyces* spp.).

ules (200 mg) morning and evening for 1 day. Pityriasis versicolor: 2 capsules (200 mg) once daily for 7 days.

- Tinea corporis, tinea cruris, tinea pedis, tinea manus:
1 capsule (100 mg) daily for 15 days; highly keratinized
regions, as in plantar tinea pedis and palmar tinea

manus, require 1 capsule (100 mg) daily for 30 days.

advisable not to give this drug to patients with a known history of liver disease. **Nursing mothers:** It is recommended not to breast feed whilst taking Sporanox (itraconazole). **Drug interactions:** Sporanox (itraconazole) should not be given concomitantly with rifampicin.

Full address: 1000 University Street, Seattle, Washington 98101

BYN OUR W AYS

**SEND FOR YOUR
FREE SAMPLE COPY**

Journal of Medical Ethics

Journal of the Institute of Medical Ethics

EDITOR: RAANAN GILLON

THE JOURNAL OF MEDICAL ETHICS

is a leading international journal in the field of medical ethics. It publishes interdisciplinary articles on ethical aspects of health care. Contributions to the journal originate in many countries and disciplines, and all undergo rigorous assessment. In addition to publishing original articles it provides a forum for reasoned debate on ethical issues. Case conferences, editorials, book reviews, correspondence and news and notes are also included. Occasional series focus on teaching medical ethics, experiences of medico-moral dilemmas ("At the Coalface") and medical ethics in literature.

RECENT AND FORTHCOMING ARTICLES:

- **The reversibility of death**
David Cole with a reply from
David Lamb
- **Debate-Involving patients in
do not resuscitate decisions**
Erich Loewy
- **Telling the truth**
Jennifer Jackson
- **Literature and medicine**
Robin Downie
- **Cost effectiveness analysis.
Is it Ethical?**
Alan Williams
- **Paying for health**
Sir Douglas Black

Annual Subscription 1992:
£69, USA (Only): \$121

**To Order Your Subscription or Sample Copy
Please Complete the Order Form Below:**

Order Form

Card Number

MasterCard users should add the numbers appearing above their name

Expiry date

Signature _____

— (Your signature is enclosed)

Name (C)

JOURNAL OF MEDICAL ETHICS

ISSN: 0306-6800

PUBLICATION: Quarterly

**PUBLICATION: Quarterly
SUBSCRIPTION RATE: \$6.00**

**SUBSCRIPTION RATE: £69
\$121 (USA ONLY)**

Please tick

Please enter my subscription, start date _____

Please send me a sample copy

Please send me *Instructions to Authors*


I enclose a cheque for _____
(Payable to British Medical Journal)

I enclose a cheque for _____
(Payable to British Medical Journal)

I wish to pay by credit card.

100

Send orders to: **BMJ Publishing Group, BMA House, Tavistock Square, London. WC1H 9JR.** or **BMJ Publishing Group, Box No 5608, Kennebunkport, Maine 04046 (Direct Orders Only).**



UK £8.95; Abroad £10.50
BMA members: £8.45 or £10.00
including postage by air abroad

How is no longer a problem — but there's still which and why

The microcomputer revolution has made powerful machines and highly complex programs generally available. This means that users of statistical techniques need no longer be concerned with the arithmetical and algebraic details — the software will take care of all that. What is vital, however, is to understand the ideas and the basic principles of statistical analysis. In *Medical Statistics on Microcomputers* R A Brown and J Swanson Beck show how to get the best use out of microcomputers when analysing data, particularly in the pathology laboratory. They explain the rational basis of various widely applicable statistical methods and also indicate their limitations so that you can make an informed choice. Chapters include:


- Data handling
- Analysis of data from one or two groups
- Comparison of several groups
- Analysis of categorical data
- Statistical methods for diagnostic tests.

Available from: BRITISH MEDICAL JOURNAL, PO BOX 295, LONDON WC1H 9TE, any leading bookseller or the BMJ/BMA bookshop in BMA House.

Please enclose payment with order, or send us full details of your MASTERCARD, VISA or AMERICAN EXPRESS credit card.

BMJ BOOKSHOP

HAS MOVED



The BMJ Bookshop has moved to larger premises at the rear of BMA House, enabling us to greatly expand the range and depth of titles we stock to cover all areas of the medical profession and related fields. We also provide a comprehensive mail order service (postage free within the UK) for customers who cannot visit the shop in person.

For further information, please telephone or write with details of books required to the address below:

BMJ Bookshop

Burton Street,
London WC1H 9JR

Tel: 071 383 6244/6638

Fax: 071 383 6662

● Euston, Kings Cross, Russell Square

Open 9.30 am. to 5.00 pm. Monday to Friday



CIA software—well written and “user-friendly”

Confidence Interval Analysis (CIA) is a computer program that takes the sweat out of calculating confidence intervals. Devised by Martin Gardner, professor of medical statistics, MRC Environmental Epidemiology Unit, University of Southampton. The program is menu driven with easy access to each chapter and to the method required within each chapter. Topics covered include calculating confidence intervals for:

- means and their differences
- proportions and their differences
- regression and correlation
- relative risks and odds ratios
- survival analyses
- non-parametric analyses


For each method relevant intermediate statistics and the required confidence interval are produced on the screen. Complete with its own manual, the program may also be used in conjunction with the book *Statistics with Confidence* (see below), which provides numerous worked examples. The software is available for IBM compatibles on either a 5½" or a 3½" disk.

“... well written, “user-friendly” ... an invaluable companion to the book and recommended highly.”
British Journal of Anaesthesia

“... highly recommended to doctors (and statisticians). CIA ... will save a lot of time and pain.”
British Medical Journal

“... an efficient and precisely focused piece of software.”
Applied Statistics

Price: £65.00. Educational establishments, research institutes, and the NHS—£45.95. Prices include VAT in the UK and air mail despatch abroad.



Over 10,000 copies sold worldwide

“... a welcome addition to the bookshelves ... as it is very practical in its approach ...”
The Pharmaceutical Journal

“The great value of this book is that it presents, in an easily understood form, the methods which will be needed in most situations.”
Journal of Epidemiology and Community Health

“... I found this book admirably suited to its purpose.”
Clinical Practice and Physiological Measurement

“This book is strongly recommended for all those who are involved in medical research and the preparation of papers for publication.”
The Medical Journal of Australia

UK £7.95; Abroad £9.50 (BMA members £7.45 or £9.00) including postage, by air abroad



BRITISH MEDICAL JOURNAL, PO Box 295, London WC1H 9TE, any leading bookseller, or the BMJ bookshop in BMA House.

Please send me _____ copy/ies of *Confidence Interval Analysis (CIA)*

£65.00 (full price) £45.95 (educational establishments, research institutes, and the NHS).

Size of disk required 5½" disk 3½" disk (please tick relevant box)

Please send me _____ copy/ies of *Statistics with Confidence - Confidence intervals and statistical guidelines*

I enclose _____ (please make cheques payable to **British Medical Journal**)

Debit my credit card American Express Mastercard Visa Card No. _____

Card Expiry Date _____ Signature _____

BMA Membership No. (where applicable) _____

NAME (BLOCK LETTERS PLEASE) _____

ADDRESS _____

POSTCODE _____

TRADEMARK

Prepulsid

Gets stomach
and oesophagus
back to work.

Prepulsid. The force behind G.I. motility.

Prescribing Information - Prepulsid (cisapride) is a gastro-intestinal prokinetic agent. Prepulsid enhances and co-ordinates gastro-intestinal propulsive motility, thereby preventing stasis and reflux. **Therapeutic indications:** 1. Gastroparesis. 2. Symptoms of X-ray or endoscopy negative upper digestive discomfort. 3. Gastro-oesophageal reflux disorders, including oesophagitis. 4. Intestinal pseudo-obstruction. **Contra-indications:** No absolute contra-indications are known. **Warnings:** Caution should be observed in patients in whom an increase in gastro-intestinal motility could be harmful. **Precautions:** *Pregnancy:* Although, in animals, there is no effect on primary fertility, no primary embryotoxic and no teratogenic effect, the anticipated therapeutic benefits should be weighed against the potential hazards before Prepulsid is given during pregnancy, especially during the first trimester. *Nursing mothers:* Although the excretion in breast milk is minimal, nursing mothers are advised not to breast feed while taking Prepulsid. **Driving and machine-operating ability:** Prepulsid does not affect psychomotor function and does not induce sedation or drowsiness. Prepulsid may, however, accelerate the absorption of central nervous system depressants, such as barbiturates and alcohol. Caution should therefore be exercised when Prepulsid is administered with these drugs. **Interactions:** - The acceleration by Prepulsid of gastric emptying may affect the rate of absorption of drugs: absorption of drugs

from the stomach may be diminished, whereas absorption of drugs from the small bowel may be accelerated (e.g. benzodiazepines, anticoagulants, paracetamol, H₂-blockers). - In patients receiving anticoagulants, the coagulation times may somewhat increase. It is advisable to check the coagulation time one week after the start of Prepulsid treatment to adapt the anticoagulant dose if necessary. The sedative effects of benzodiazepines and of alcohol may be accelerated. - The effects of Prepulsid on gastro-intestinal motility are, for the most part, antagonized by anticholinergic drugs. - In hepatic and renal insufficiency, it is recommended to halve the initial daily dose. Subsequently, this dose can be adapted, depending on the therapeutic effects or possible side-effects. - In the elderly, steady-state plasma levels are generally higher, due to a moderate prolongation of the elimination half-life. Therapeutic doses, however, are similar to those used in younger patients. - In the case of drugs that require individual titration, it may be useful to monitor plasma levels of such drugs when Prepulsid is associated. **Adverse reactions:** In line with the pharmacological activity of Prepulsid, transient abdominal cramping, borborygmi and diarrhoea may occur. Mild and transient headache or lightheadedness have been reported occasionally. When diarrhoea occurs in babies or infants, the dose should be reduced. There are isolated reports of CNS effects, i.e. convulsive seizures and

extrapyramidal effects. **Dosage:** - Adults: according to the severity of the condition, 15 to 40 mg daily, to be given in 2 to 4 intakes, to be taken as tablets or as oral suspension (the full plastic 5-ml spoon contains 5 mg). As a rule the following doses have proven adequate: • less severe conditions: 5 mg t.i.d. (dose can be doubled); • severe conditions (gastroparesis, oesophagitis, refractory constipation): 10 mg t.i.d. to 10 mg q.i.d. (before the 3 main meals and before retiring). - Infants and children: on the average 0.2 mg/kg per intake, 3 to 4 times daily. For the suspension, intakes are indicated on the dosing pipet as a function of body weight.

Full prescribing information available on request.

Note: Prepulsid (cisapride) is not yet available in all countries and not all indications have been approved everywhere.

 **JANSSEN**
PHARMACEUTICA
B-2340 Beerse, Belgium