


The

British Medical Journal



THE JOURNAL OF THE BRITISH MEDICAL ASSOCIATION.

Including an Epitome of Current Medical Literature.

WITH SUPPLEMENT.

No. 3347.

SATURDAY, FEBRUARY 21, 1925.

Price 1/3

Influenza Prophylaxis

IT has been established that the serious symptoms of epidemic influenza are due to infection by the streptococcus, pneumococcus and influenza bacillus, and experience has proved that a mixed vaccine containing these organisms is of great value as a prophylactic agent. The proportions considered most suitable for the purpose are provided in the MIXED ANTI-INFLUENZA VACCINE (St. Mary's Hospital Formula).

For those who have had repeated attacks of epidemic influenza, it may be preferable to employ the more comprehensive ANTI-CATARRH VACCINE (Prophylactic) which was particularly designed for immunization against ordinary nasal and bronchial catarrhs.

A booklet, "Vaccine Therapy," containing full particulars, will be sent post free on request by

PARKE, DAVIS & Co., LONDON, W.1.

Telegrams:
"CASCARA, LONDON."

Telephone No.:
REGENT 3260.

Mixed Anti-Influenza Vaccine

(St. Mary's Hospital Formula)

Each c.c. contains:

B. influenzae.....500 millions
Pneumococcus (*mixed types*) 1000 millions
Streptococcus (*mixed types*)...100 millions

Anti-Catarrh Vaccine (Prophylactic)

Each c.c. contains:

M. catarrhalis.....50 millions
B. pneumoniae.....50 millions
Pneumococcus.....250 millions
B. septus.....50 millions
B. influenzae.....250 millions
Streptococcus.....25 millions

The above-mentioned vaccines are supplied in sets of three graduated doses for the immunization of one adult; also in ampoules of 1 c.c., and in bottles of 25 c.c.

These vaccines are prepared in the laboratory of the Department for Therapeutic Inoculation, St. Mary's Hospital, London (Director: Sir Almroth Wright, M.D.).