


DEC 1 1923

The Medical Lib.

# British Medical Journal



THE JOURNAL OF THE BRITISH MEDICAL ASSOCIATION.

*Including an Epitome of Current Medical Literature.*

WITH SUPPLEMENT.

No. 3281.

SATURDAY, NOVEMBER 17, 1923.

Price 1/3

## PHYLACOGENS

Phylacogens are modified bacterial derivatives manufactured by Parke, Davis & Co. Their composition is based on the theory of mixed infections, and they represent the end products of cell digestion. They produce a rapid immunising response and may be administered intravenously or hypodermically.

**ERYSIPELAS PHYLACOGEN** is indicated in the treatment of erysipelas and other acute infectious diseases caused by streptococci.

**GONORRHOEA PHYLACOGEN.** Excellent results are reported from the hypodermic administration of Gonorrhoea Phylacogen in chronic pathological conditions due to infection by *M. gonorrhoeæ*.

**PNEUMONIA PHYLACOGEN.** Indicated in the treatment of pneumonia, or any pathological condition where *Diplococcus pneumoniae* is the primary etiological factor.

**MIXED INFECTION PHYLACOGEN.** Indicated in all infections, acute or chronic, mild or severe, in which no specific micro-organism is found to be predominant, notably in surgical infections, abscesses, puerperal sepsis, etc.

**RHEUMATISM PHYLACOGEN.** Has proved remarkably effective in a number of rheumatic conditions where streptococci constitute an important factor in the etiology.

**TYPHOID PHYLACOGEN.** Indicated in the treatment of typhoid fever or any pathological condition caused by infection with *B. typhosus*.

*Phylacogens are supplied in 1 c.c. ampoules and 10 c.c. phials. For further particulars please refer to our booklet "Phylacogen Therapy," which will be sent to physicians, post free, on request.*

**PARKE, DAVIS & CO., 50, Beak St., Regent St., LONDON, W.1.**

Telephone: REGENT 3260 (six lines).

Telegrams: "CASCARA," LONDON.