


DEC 20 1926

Medical Lib.

The

British Medical Journal

THE JOURNAL OF THE BRITISH MEDICAL ASSOCIATION.



Including an Epitome of Current Medical Literature.
WITH SUPPLEMENT.

No. 3433.

SATURDAY, DECEMBER 4, 1926.

Price 1/3

Scarlet Fever Antitoxin

SCARLET FEVER STREPTOCOCCUS ANTITOXIN, CONCENTRATED (Globulin)

(Parke, Davis & Co.)

- Q For Treatment.** A dose of 10 c.c. of this antitoxin usually establishes normal temperature within 36 hours, and the quarantine period is shortened in many cases by nearly 50 per cent. Early administration generally arrests the progress of complications, and in some cases has mitigated the severity of complications already developed.
- Q For Prophylaxis.** Results so far reported seem to indicate that a dose of 2.5 c.c. of this antitoxin administered to children of from 5 to 15 years of age will confer passive immunity for from 10 to 14 days.
- Q For Diagnosis.** The *diluted* antitoxin injected intradermally into an area where the suspected rash is most marked will, in a case of scarlet fever, produce the diagnostic rash-fading reaction of Schultz-Charlton.

Further particulars will be sent on request.

PARKE, DAVIS & COMPANY



50 BEAK ST., LONDON, W.1

Telephone: Regent 7801

Telegrams: "Cascara, London"