


The

DEC 1 1927

Medical Library

British Medical Journal

THE JOURNAL OF THE BRITISH MEDICAL ASSOCIATION.



**Including an Epitome of Current Medical Literature
WITH SUPPLEMENT.**

No. 3489.

SATURDAY, NOVEMBER 19, 1927.

Price 1/3

For Amylaceous Dyspepsia

PARKE, DAVIS & Co., introduced Taka-Diastase to the medical profession more than a quarter of a century ago, and it was not long before it came to be recognised as the standard remedy for so-called amylaceous dyspepsia.

Without question, Taka-Diastase is the most powerful of all amylolytic agents of vegetable origin. It is standardised to render soluble three hundred times its weight of dry starch in ten minutes, the starch being transformed to dextrins, maltose and dextrose.

Taka-Diastase is also an active proteolytic and, further, acts in neutral, slightly alkaline or slightly acid media. Whilst a very active therapeutic agent, Taka-Diastase is perfectly harmless. Being tasteless, it is freely taken by children and fastidious patients.

Taka-Diastase

Supplied as powder, tablets or liquid. Tablets are also available containing Taka-Diastase with pepsin, pancreatin or strychnine. It may also be obtained as an elixir, with bismuth, nux vomica and hydrocyanic acid. Further particulars will be sent to any member of the medical profession on request.

D

PARKE, DAVIS & COMPANY



50 BEAK ST., LONDON, W.1.