

The

British Medical Journal

THE JOURNAL OF THE BRITISH MEDICAL ASSOCIATION



Including an Epitome of Current Medical Literature
WITH SUPPLEMENT

No. 3737

SATURDAY, AUGUST 20, 1932

Price 1/3

PERCAINE "CIBA"

Trade Mark Registered.

The New Local Anaesthetic for Regional, Infiltration, Surface and Spinal Anaesthesia

Acts in extreme dilution (0.5-2: 1000).

Produces anaesthesia of hitherto unattained intensity and duration.

Not a narcotic.

Economical in use.

Belongs chemically to a class entirely different from cocaine and its derivatives.

Vide *British Medical Journal*, March 15, 1930, pp. 488-9 and 495-6, and April 5, 1930, pp. 669-70. *The Lancet*, March 15, 1930, pp. 573-4 and 587. *British Journal of Anaesthesia*, April and July, 1930, and January, 1931. *Proceedings of the Royal Society of Medicine*, May, 1930, pp. 919-928. *British Journal of Urology*, June, 1930, pp. 129, 130, and 179. *Edinburgh Medical Journal*, April, 1931. *St. Bartholomew's Hospital Journal*, February, 1931, pp. 93-96, etc., etc.

Packages available:

Percaine (Hydrochloride) Crystals.
(For solutions) 5 grm.

Percaine Base.
(For ointments and oily solutions) 5 grm.

Percaine Tablets (for the preparation of solutions).
Tubes of 20 × 0.05 grm. Tubes of 10 × 0.1 grm.

Percaine-Adrenalin Tablets.
Tubes of 10.

Percaine Solution 2% (exclusively for surface anaesthesia in E.N.T. work). Stoppered bottles of 30 c.c.

Percaine Ampoules.

Boxes of 5 × 5.5 c.c. Solution 1: 1000 (with Adrenalin).
Boxes of 2 × 30.5 c.c. Solution 1: 1000 (with Adrenalin).
Boxes of 10 × 30.5 c.c. Solution 1: 1000 (with Adrenalin).

Boxes of 10 × 2.3 c.c. Solution 2: 1000 (with Adrenalin).
Boxes of 10 × 2.3 c.c. Solution 3: 1000 (with Adrenalin).
Boxes of 12 × 20 c.c. Solution 1: 1500, for Spinal Anaesthesia.
Boxes of 10 × 2.3 c.c. Buffered Solution 1: 200, for Spinal Anaesthesia.

THE CLAYTON ANILINE Co. Ltd., 40 SOUTHWARK STREET, LONDON, S.E.1

Telephones: Hop 4474 (3 lines).

Pharmaceutical Department.

Telegrams: Cibadyes Boroh London