448.8




JOURNAL OF THE BRITISH MEDICAL ASSOCIATION

> JULY 4 1 1959 A RECORD SATURDAY

CONTENTS

U. S. DEPARTMENT OF ADDICULTURES An Experiment in Mass Radiography J. B. COCHRANE, M.B., M.R.C.P., W. B. FLETCHER, F.S.S., and C. CLAYSON, M.D. **Puerperal Cardiomyopathy** S. M. Rosen, M.B. **Acute Bacillary Dysentery in Cyprus** P. J. TAYLOR, M.B., M.R.C.P. Sulphonamides and Streptomycin in Bacillary Dysentery 



The advantages of

## Esidrex

JUL 30 1959

(hydrochlorothiazide, a product of CIBA research)

## For oedema and hypertension

Highest potency-low dosage.

Fully active orally-often effective where mercurials have failed.

Less risk of potassium depletion.

Almost a pure sali-diuretic.

No significant effect on urinary pH.

Low cost of treatment—averages 4d. to 1/- daily.

The dosage range is usually 25 to 100 mg. daily, taken in two doses, after breakfast and after lunch.

Tablets of 25 mg. and 50 mg.\* hydrochlorothiazide in 25's, 100's and 500's.

\*Available shortly.

Esidrex is a registered trade mark. Reg. user

CIBA Laboratories Limited, Horsham, Sussex.

BRITISH MEDICAL ASSOCIATION, TAVISTOCK SQUARE, LONDON, W.C.1.

No. 5139 WEEKLY, PRICE 3s.

2/9

Copyright 1959

REGISTERED AS A NEWSPAPER