



SATURDAY 24 FEBRUARY 1973

LEADING ARTICLES

Symptoms in Hypertension page 433 Care of Elderly People with Dementia page 434 B.C.G. Tested page 435 Not So Popular Psychiatry page 435 "Seborrhoeic Dermatitis" Problems of Iron-deficiency Anaemia in Infancy page 437 page 436 "Steal" Syndromes and Hypertension page 438

PAPERS AND ORIGINALS

Cryptogenic Drop Attacks: An Affliction of Women D. L. STEVENS, W. B. MATTHEWS.	439
Plasma Digoxin Concentration in Children with Heart Failure J. E. CREE, D. J. COLTART, M. R. HOWARD	443
Rubella Epidemic in a Maternity Unit STUART CARNE, C. J. DEWHURST, ROSALINDE HURLEY	
Effect of Metoclopramide on Gastric Emptying during Labour F. A. HOWARD, D. S. SHARP	
Effect of Oestrogen Therapy on Plasma and Urinary Levels of Uric Acid ANNE NICHOLLS, M. L. SNAITH, J. T. SCOTT	
Removal of Polyps with Fibreoptic Colonoscope: A New Approach to Colonic Polypectomy	
CHRISTOPHER WILLIAMS, T. MUTO, K. R. P. RUTTER	451
Operative Colonoscopy H. J. ESPINER, P. R. SALMON, R. H. TEAGUE, A. E. READ	453
Nature of Hyperacute (Accelerated Second Set) Rejection in Dog Renal Allographs and Effects of Heparin on	
Rejection Process A. H. AMERY, G. D. PEGRUM, R. A. RISDON, GRANT WILLIAMS	455
Amodiaquine-induced Agranulocytosis: Toxic Effect of Amodiaquine in Bone Marrow Cultures In Vitro	
D. E. LIND, J. A. LEVI, P. C. VINCENT	
Lymphosarcoma Presenting as Löffler's Syndrome C. C. BAILEY, R. H. A. CAMPBELL	460
Severe Cardiovascular Complication Associated with Prolonged Starvation	
F. SANDHOFER, F. DIENSTL, K. BOLZANO, H. SCHWINGSHACKL	
Spontaneous Oesophageal Rupture after Carbachol Therapy P. COCHRANE	463
Erysipelothrix Septicaemia without Endocarditis R. H. TOWNSHEND, A. E. JEPHCOTT, M. HASAN YEKTA	464

MEDICAL PRACTICE

Review of Schizophrenics in a Rural Practice over 26 Years	C. A. H. WATTS
Extended Deep Femoral Angioplasty R. BERGUER, L. T. COTTON	, S. SABRI
Large-bowel Perforations in Patients Undergoing Sigmoidosc	
Role of Teaching Hospitals in a Developing Country 0. 0. AF	KINKUGBE
A New Look at Infectious Diseases: Hepatitis G. C. TURNER	
Any Questions?	
Personal View CLIFFORD HAWKINS	
CORRESPONDENCE—List of Contents	OBITUARY NOTICES

NEWS AND NOTES

Epidemiology—Orf and Milkers' Nodes	497
Medicolegal—Seat Belts and Negligence	497
Parliament-Changes to N.H.S. Reorganization Bill	498
Medical News—G.M.C. Inquiry Committee	499

ODITO		NOTIOLS	••••	• • • •	••••	••••	•••••	••••••
BOOK	RE	VIEWS					••••	496

SUPPLEMENT

Special Conference of Representatives of Local Medical Committees—Chambers Report and Repre-	
sentation of N.H.S. General Practitioners	51
Association Notices	60

CORRESPONDENCE

An Integrated Child Health Service

Correspondents are asked to be brief

Hyperactive Children G. F. J. Goddard, D.P.M.; R. Payne,
F.R.C.PSYCH
Influenza Outbreak
S. K. Mandal, M.R.C.P481
Certification of Hypothermia Deaths
A. M. Adelstein, M.D
Hypothermia
R. Wigglesworth, F.R.C.P
Nomenclature of the New Vagotomy
C. Wastell, F.R.C.S., and others
Gastric Ulcer after Highly Selective
Vagotomy
D. Johnston, F.R.C.S.ED
"Twin" Intracranial Aneurysms
P. J. E. Wilson, F.R.C.S., and I. P. Cast,
F.R.C.S.; W. M. Chesney, M.R.C.G.P
Gonococcaemia in the Family
J. K. Oates, F.R.C.P.ED., and V. H. Patterson,
B.CHIR
How Infectious is Gonorrhoea?
M. E. Ward, PH.D., and P. J. Watt, M.D485
Glove-powder Peritonitis
D. F. E. Nash, F.R.C.S
Pulmonary Oedema in Pulmonary
Thromboembolism
W. J. Windebank, M.R.C.P., and F. Moran,
F.R.C.P.ED
Pathology of Malignant Hyperpyrexia
W. G. Bradley, D.M
Thiocyanate Metabolism in Human Vita-
min B ₁₂ Deficiency
D. McAlpine, F.R.C.P
Medical Aspects of Ambulance Design
R. N. Seymour, M.B486

Staffing of Accident and Emergency
Departments
R. K. Holdsworth, B.M.; J. P. Turney,
F.R.C.S
Disappearance of Diphtheria
M. G. McEntegart, M.D., F.R.C.PATH., and
I. Zamiri, м.в
Dyslexia as Cause of Psychiatric Disorder
Beve Hornsby, M.Sc
Treatment of Malaria
W. Peters, м.D
Complication of Internal Jugular Vene-
puncture
G. A. Matthews, F.R.C.PATH
Possible Hazard of Contact Lens Manu-
facture
R. Routledge, D.I.H
Making Hospital Geriatrics Work
Margaret B. Hawker
Sickle-cell Anaemia
Colonel H. Gall, M.R.C.S488
Nitrazepam: Another Interesting Syn-
drome
Н. MacLean, м.в
Internal Fixation of Secondary Deposits in
Long Bones
W. H. Bond, F.R.C.S., D.M.R.T., and others488
Anthrax
M. H. Hughes, F.R.C.PATH488
Glucagon and Growth Hormone
L. Stimmler, м.D489
Ethics in Epidemiological Research
S. Bradshaw, M.B

Helenor F. Pratt, M.B., D.C.H.; A. J. Essex-W. K. S. Moore, M.B.; T. G. F. Hudson, F.R.C.P. . . Children's Wheelchairs Sponsored Hospitality **Insulin Syringes** Fees for Cremation Certificates D. A. Spencer, M.B., M.R.C.PSYCH.491 Side Effects of Lithium Carbonate D. P. Richardson, M.SC., and M. Lawson, Anomalies in Tendon Reflexes Lieutenant-Colonel V. B. Whittaker, G. Bousfield, M.D......491 Infantile Mumps B. S. Milner, M.B......491 Chondromalacia Patellae J. Darracott, D.PHYS.MED., and B. Vernon-Screening for Cervical Carcinoma Management of the N.H.S. D. G. Ferriman, F.R.C.P......492 Ten-session Consultant Contract L. J. Beilin, M.D., and others......492

Hyperactive Children

SIR,-Your leading article on what you call "hyperactive" children (10 February, p. 305) took much too wide a view, and the very well-recognized and well-recognizable syndrome of hyperkinesis got completely hidden from sight behind a welter of side issues. The only psychopathological disturbance in hyperkinesis is overactivity, and there is only one cause of overactivity and that is hyperkinesis. Domestic and educational stresses do not cause hyperkinesis any more than does parental alcoholism or criminality, parent-child conflicts and sibling rivalry (which are universal problems anyway), repression at school (which is a necessity), or parental drug addiction. Hyperkinesis can certainly leave all these things in its wake.

What alarms me is the suggested treatment of hyperkinesis, in which, I agree, drugs have an important role. The use of methyl phenidate and, more important, of dexamphetamine is fraught with the danger of addiction, which is a certainty if the upper limit of age is accepted as 16 years. Any child who continues to be called "hyperkinetic" after the age of 12 is not suffering from the hyperkinetic syndrome, and the use of stimulant drugs is contraindicated. Diphenhydramine is an antihistamine, one of the side effects of which is drowsiness, and soporifics in general are not only useless in treatment but can be positively dangerous by increasing the accident-proneness which already exists in hyperkinesis. The phenothiazines are sometimes symptomatically useful, but

medication with such drugs even for a relatively short period leads to obesity, which is possibly more incapacitating. The most effective drug in my experience is the butyrophenone haloperidol, which is avail-able in a drop bottle, one drop containing 0.1 mg of the drug. Effective dosage can be reached by trial and error and the safety margin is very large. A starting dose of one drop twice daily can be increased, according to response, by one or two drops each dose, the average effective dose being between four and 15 drops daily. Adverse side effects are rare and can be reversed by reducing the dosage without impairing the symptomatic response, and there is no recorded case of addiction.-I am, etc.,

Halifax, Yorks.

G. F. J. GODDARD

SIR,—In your leading article (10 February, p. 305) you state that minimal cerebral dysfunction is one of the causes of hyperactivity in children.

Cerebral function is either normal or abnormal, and no child should be branded as having a dysfunctioning brain, however minimal, unless he suffers from neurological disorder or mental handicap.—I am, etc.,

RUDOLPH PAYNE I am, etc.,

Little Plumstead Hospital, Norwich

Influenza Outbreak

SIR,—I was most interested to read the letter from Dr. C. S. Goodwin and others (13 January, p. 109), in which they demonstrated clearly the effectiveness of influenza vaccine among geriatric patients. I have also observed the same sort of feature in the recent outbreak.

In the Border group of hospitals in Wales the geriatric wards for continuing care and rehabilitation are scattered in the peripheral hospitals. In one such ward, at Bronllys Hospital, where patients were vaccinated (with Admune which, like Influvac, does not contain influenza virus A/Eng/42/72), none of the 15 patients suffered from influenza, whereas in other wards where patients were not vaccinated 14 of the 25 patients at Llandrindod Wells Hospital developed influenza and one died, and 15 of the 46 patients at St. David's Hospital, Brecon, developed influenza and four died. These facts obviously suggest that elderly patients do benefit from influenza vaccine.

However, I believe that there is still a great deal of doubt about the wisdom of vaccinating elderly patients in a continuing care (long-stay) type of ward, even though people generally agree that geriatric patients in acute and rehabilitation wards should be vaccinated.

I would welcome the views of the medical profession at large on this particular issue.— I am, etc.,

S. K. MANDAL St. Tydfil's Hospital, Merthyr Tydfil, Glam.